

How valuable is physical examination of the cardiovascular system?

Andrew Elder,¹ Alan Japp,² Abraham Verghese³

¹Department of Geriatric Medicine, Edinburgh Medical School, Edinburgh, UK

²Department of Cardiology, Edinburgh Medical School, Edinburgh, UK

³Department of Medicine, Stanford University, Palo Alto, CA, USA

Correspondence to: A T Elder, Department of Acute Medicine of Older Age, Western General Hospital, Edinburgh EH4 2XU, UK
atelder@gmail.com

Cite this as: *BMJ* 2016;354:i3309
doi: 10.1136/bmj.i3309

ABSTRACT

Physical examination of the cardiovascular system is central to contemporary teaching and practice in clinical medicine. Evidence about its value focuses on its diagnostic accuracy and varies widely in methodological quality and statistical power. This makes collation, analysis, and understanding of results difficult and limits their application to daily clinical practice. Specific factors affecting interpretation and clinical application include poor standardisation of observers' technique and training, the study of single signs rather than multiple signs or signs in combination with symptoms, and the tendency to compare physical examination directly with technological aids to diagnosis rather than explore diagnostic strategies that combine both. Other potential aspects of the value of physical examination, such as cost effectiveness or patients' perceptions, are poorly studied. This review summarises the evidence for the clinical value of physical examination of the cardiovascular system. The best was judged to relate to the detection and evaluation of valvular heart disease, the diagnosis and treatment of heart failure, the jugular venous pulse in the assessment of central venous pressure, and the detection of atrial fibrillation, peripheral arterial disease, impaired perfusion, and aortic and carotid disease. Although technological aids to diagnosis are likely to become even more widely available at the point of care, the evidence suggests that further research into the value of physical examination of the cardiovascular system is needed, particularly in low resource settings and as a potential means of limiting inappropriate overuse of technological aids to diagnosis.

Introduction

Some of the earliest accounts of the interaction between a patient and his or her doctor concern the physical examination of the cardiovascular system.¹ The primary purpose of such examination is to assess the presence, nature, and severity of cardiovascular disease. Several specific components of the physical examination have been described to support these aims.

Substantial evidence of the diagnostic value of specific elements of the physical examination of the cardiovascular system exists,^{2,3} but its methodological quality varies widely. Since the advent of technological aids to diagnosis (TAD) such as echocardiography in the late 1970s,⁴ the central role of physical examination in the practice of clinical cardiology has been challenged,⁵⁻⁷ and its standard of practice has declined in some countries.⁸⁻¹²

Despite this, many clinicians still use and attach significance to physical examination techniques in daily practice. In acute general internal medicine, the combination of history and physical examination without recourse to TAD still accounts for most diagnoses.¹³ Findings from the physical examination of the cardiovascular system

are central to diagnostic scoring systems for conditions such as pulmonary embolism,¹⁴ risk stratification in acute coronary syndromes,¹⁵ and "early warning scores" in patients admitted to hospital.¹⁶ In medical education, the physical examination is still regarded as a core component of undergraduate and postgraduate curriculums.

This review will summarise the evidence relevant to the value of the physical examination of the cardiovascular system in contemporary clinical practice, comment on the quality and limitations of the evidence, and highlight areas where further investigation may be most appropriate. It is aimed at clinicians interested in the evidence base for the physical examination they undertake in daily clinical practice and those who teach doctors in training. It is also relevant to people with a research interest in diagnostic methods, the doctor-patient relationship, and over-investigation and overuse of technology in healthcare.

Sources and selection criteria

The physical examination of the cardiovascular system is traditionally divided into several components, summarised in figure 1.

Component	Specific features of interest
Palpation of arterial pulse	<ul style="list-style-type: none"> • Rate • Rhythm • Volume • Character or contour • Vessel wall • Bruits • Symmetry
Measurement of arterial blood pressure	<ul style="list-style-type: none"> • Systolic pressure • Diastolic pressure • Pulse pressure
Examination of jugular veins	<ul style="list-style-type: none"> • Venous pressure • Venous waveform
PRAECORDIAL EXAMINATION:	
Inspection	<ul style="list-style-type: none"> • Scars • Abnormal movements
Palpation	<ul style="list-style-type: none"> • Apical impulse (location and character) • Praecordial impulses • Thrills
Percussion	<ul style="list-style-type: none"> • Cardiac dullness
Auscultation	<ul style="list-style-type: none"> • Heart sounds (normal and added) • Murmurs • Pericardial rub
General and other systems examination	<ul style="list-style-type: none"> • Cyanosis • Clubbing • Oedema • Cutaneous features of peripheral arterial or venous disease • Peripheral signs of infective endocarditis • Transverse earlobe crease • Arcus lipidus • Xanthelasma and xanthoma • Calf tenderness and swelling • Capillary refill time • Pulmonary crepitations (rales) • Hepatomegaly • Splenomegaly • Funduscopy

Fig 1 | Components of physical examination of the cardiovascular system

The purpose of each component of the physical examination is to identify physical signs, defined here as a variation from an expected or normal clinical observation or finding. As the evidence base on physical examination of the cardiovascular system primarily focuses on the diagnostic accuracy of physical signs, this review also focuses on diagnostic accuracy but acknowledges that the value of physical examination may extend beyond this single parameter.

We have not:

- Provided a detailed account of the physiology underlying normal findings in the physical examination or the pathophysiology causing specific physical signs suggestive of cardiovascular disease.
- Reviewed the value of any aspect of the physical examination in which a device other than a stethoscope or a sphygmomanometer is used.
- Considered paediatric practice or the evaluation of congenital heart disease in adults.

- Reviewed the prognostic value of measurements of systolic or diastolic blood pressure, their mean, or their difference (pulse pressure), as their use in cardiovascular risk assessment and treatment is established.
- Reviewed established diagnostic scoring or risk stratification systems, or early warning scores that include components of the physical examination—for example, those used in the diagnosis of pulmonary embolism,¹⁴ deep venous thrombosis,¹⁷ infective endocarditis,¹⁸ or chronic venous insufficiency.¹⁹

Although we did an extensive literature review, we did not apply specific methods demanded of formal systematic reviews or meta-analyses. We used an electronic search of PubMed based on a strategy recommended in relation to the diagnostic value of physical examination to identify studies published between May 1965 and May 2015.²⁰ This combined terms relating to clinical entities (AND)* terms relating to clinical skills (AND)* terms relating to diagnostic testing. Figure 2 shows the terms used.

We also identified and reviewed all references in the three existing major contemporary sources of collated evidence on physical examination,²⁻²¹ as well as publications citing these references since their publication date. We included clinical practice guidelines, but physical examination is not always included in such guidelines and often not evaluated in similar detail to data from other sources. We reviewed only English language publications, and we did not consider further any articles that established pathophysiological mechanisms underlying physical signs, single case reports of associations between signs and specific diseases, and articles that provided no numerical data to enable estimation of diagnostic accuracy.

The primary electronic search strategy failed to identify approximately 20% of the publications found in the three collated sources, their references, or their citations. The difficulty of identifying publications relating to the diagnostic value of physical examination has been previously identified.²²

Interpreting the literature

The following factors should be considered when interpreting the literature on the value of physical examination of the cardiovascular system.

Definition of value

In the context of physical examination, the word “value” is generally taken to be synonymous with the diagnostic accuracy of physical signs when specific diagnoses are being considered—that is, whether the presence or absence of a sign or combination of signs makes a diagnosis more or less likely. Most studies of physical examination are based on this construct and typically compare the physical examination with a diagnostic reference standard, usually an imaging or invasive haemodynamic study.

This definition may underestimate the true value of physical examination, and a broader definition could include the factors shown in figure 3. The literature contains little information on factors other than diagnostic accuracy and contributions to clinical care beyond diagnosis.




Fig 2 | Search terms used in electronic search

Methodological considerations

Ideal standards for tests of diagnostic accuracy have been described, but many studies of the diagnostic accuracy of physical examination fail to meet such criteria.²² For example, studies are generally small; participants are often poorly characterised beyond age, sex, and primary diagnosis; analyses are typically univariate rather than multivariate; and there are relatively few systematic reviews or meta-analyses. The following factors must also be considered.

Characteristics of clinician observers and observer variability

In studies of the diagnostic accuracy of physical examination, observations provided by clinicians provide the data on which analysis of diagnostic value is based. The number of clinician observers is highly variable—typically less than six—and some studies include only one observer. Studies in which a single clinician does all observations show that high levels of accuracy and low levels of intra-observer variability for some signs can be obtained.²⁷⁻²⁹

Most studies include more than one clinician observer and describe varying degrees of inter-observer variability but provide no information to indicate how the observers' technique of physical examination was assessed, standardised, or compared and characterise observers only according to their specialty, job title, or training grade. Inter-observer variability varies according to the experience or seniority of the participating clinicians and can be reduced with training of observers.³⁰⁻³² These find-

ings suggest that observers' technique influences inter-observer variability; although the overall contribution of observers' characteristics to such variability is difficult to quantify, it is likely to be substantial.

Characteristics of clinical observations

Estimations of the diagnostic accuracy of physical examination assume that a sign is binary—that is, it is either present or absent. In many clinical situations, however, clinicians accord diagnostic weight according to the degree of positivity of a sign or their level of certainty that it is present or absent. For example, a trace of peripheral oedema may be ascribed lower diagnostic significance than oedema that is present above the knee. In general, the literature takes no account of this. In addition, the evanescent nature of some signs (such as pericardial rub) makes structured analysis of diagnostic value challenging.

Characteristics of clinical practice

The literature on physical examination of the cardiovascular system typically compares physical examination directly with TAD. In reality, clinicians use combinations of diagnostic modalities that typically include history, physical examination, blood tests, and imaging. Few studies attempt to evaluate the comparative efficacy of strategies that use combinations of diagnostic modalities (for example, physical examination plus TAD) compared with single modality (for example, physical examination alone or TAD alone) approaches. In addition,



Fig 3 | Factors contributing to the value of physical examination of the cardiovascular system. TAD=technological aids to diagnosis

combinations of signs and symptoms are typically used in clinical diagnostic and therapeutic decision making, but much of the literature attempts to evaluate single signs in isolation from other physical examination findings or ancillary clinical information such as the history. Exceptions are prediction scores for pulmonary embolism or deep venous thrombosis and some studies in heart failure diagnosis.

Global variation of disease prevalence

Most studies of the diagnostic utility of physical examination have been conducted in the United States or Europe. This may account for the lack of studies from which evidence relating to the utility of the physical examination in the diagnosis of conditions now less commonly seen in Western populations, such as rheumatic mitral stenosis, can be derived.

Diagnostic reference standards

TAD are typically used as the diagnostic reference standard in studies of physical examination, which raises two methodological considerations. Firstly, different definitions of abnormality may be applied in different studies of the same sign. For example, the ultrasound definition of a “significant” arterial stenosis varies in studies of the diagnostic value of pulse palpation or bruits. Secondly, many studies provide no information on the (intra- or

inter-)observer variability of the diagnostic reference standard, either as used in the study or in general use. Studies in which inter-observer variability is described show the presence of varying degrees of such variability in the interpretation of commonly used TAD. For example, echocardiographic assessments of various parameters of left ventricular systolic and diastolic function contain observer variability,^{33 34} which may both influence the interpretation of study results and be underappreciated by many clinicians.²

Range of statistical indices used

The diagnostic value of a test is determined by its accuracy and precision (reliability).³⁵ A wide variety of statistical indices are used to express diagnostic value in studies of physical examination of the cardiovascular system,³⁶ and comparison between or pooling of studies is challenging, particularly as the raw data necessary to calculate missing indices are often absent from the published papers.

Likelihood ratios provide information on the probability of disease in an individual³⁷; when available, they are the preferred statistical index of comparison in this review. Figure 4 shows the definition of likelihood ratios and their method of calculation, and figure 5 illustrates the relation between a likelihood ratio and diagnostic pre-test and post-test probability.

STATE OF THE ART REVIEW

Table 1 | Summary of evidence of clinical value

Sign or manoeuvre	What is this sign purported to indicate?	Suggested clinical value
Arterial pulses		
Rate	Cardiac rate	Non-specific diagnostically but has prognostic value in several acute settings. ⁴¹ Resting heart rate predicts mortality in population settings ⁴²
Postural tachycardia	Hypovolaemia	A rise of 30 beats/min on standing suggests large blood loss ⁴³
Rhythm	Cardiac rhythm	See detailed discussion of specific signs
Volume and contour	Changes in stroke volume or vascular compliance	
Slow rising/plateau	Aortic stenosis (AS)	Supports diagnosis of AS in patients with systolic murmur ²⁹
Collapsing/water-hammer	Aortic regurgitation (AR)	Insensitive and non-specific when assessed by palpation. High (>80 mm Hg) measured pulse pressure has LR+ of 10.9 for moderate to severe AR in a single study ⁴⁴
Pulsus bisferiens	Aortic valve disease (AS, AR), HCM	Low specificity and sensitivity
Dicrotic	Mitral regurgitation, high SVR, low stroke volume	Low specificity and sensitivity
Jerky	HCM	Said to be pathognomic
Small volume/weak/thready	Shock, impaired LV function, AS	Low specificity and sensitivity
Other:		
Pulsus paradoxus	Cardiac tamponade	In patients known to have a pericardial effusion, if pulsus paradoxus of >12 mm Hg is present, the LR+ for cardiac tamponade is 5.9 (95% CI 2.4 to 14.3) and if pulsus paradoxus is absent the LR- is 0.03 (0 to 0.2) ^{45,46}
Pulsus alternans	LV systolic impairment	Moderate specificity but low sensitivity ⁴⁷
31 eponymous signs of AR	Presence or severity of AR	Generally specific findings but low sensitivity ⁴⁸
Absence of specific pulses	Obstructive disease at or proximal to site of palpation	See detailed discussion of specific signs
	Hypovolaemic shock	One study suggests that a palpable femoral pulse means shock is less likely ⁴⁹
Buerger's test	Peripheral arterial disease (PAD)	May suggest more severe PAD when present ⁵⁰
Capillary refill time	Peripheral hypoperfusion	See detailed discussion of specific signs
Pulse asymmetry: upper limbs	Obstruction at specific site	
	Aortic dissection	See detailed discussion of specific signs
	Subclavian stenosis and steal syndrome	Asymmetry has high sensitivity in patients with suggestive symptoms ⁵¹
Pulse asymmetry: upper and lower limbs	Coarctation of the aorta	Combinations of blood pressure and pulse findings have high sensitivity in children ⁵²
Bruits at specific sites	Obstructive arterial disease	
Abdominal	Renovascular disease	See detailed discussion of specific signs
Carotid	Carotid artery disease	See detailed discussion of specific signs
Pulsatile abdominal mass	Abdominal aortic aneurysm	See detailed discussion of specific signs
Jugular venous pulse		
Jugular venous pressure	Central venous/right atrial pressure	See detailed discussion of specific signs
Kussmaul's sign	Constrictive pericarditis	Low sensitivity
Response to Valsalva	Elevated left heart filling pressure	LR of >5 in a single study ⁵³
Abdominojugular test (hepatojugular reflux)	Elevated left heart filling pressure	See detailed discussion of specific signs
Character of the venous waveform	Abnormalities of intracardiac blood flow or cardiac rhythm	
Giant v wave	Tricuspid regurgitation	LR+ of 10.9 in a single study ²⁹
Other abnormalities of a/c/v waves or x/y descent	Abnormalities of blood flow into heart or heart rhythm	Generally, low sensitivity and specificity
Cannon waves	Atrioventricular dissociation	In a single study of induced ventricular tachycardia, cannon waves in the jugular venous waveform had a negative LR of 0.1 ³¹
Other veins:		
Peripheral venous collapse	Elevated central venous pressure	One study suggests low sensitivity but high specificity ⁵⁴
Praecordium		
Percussion of the heart	Cardiac size	Single study suggests value in excluding cardiomegaly ⁵⁵
Palpation of apical impulse:		
Location	Cardiac size and LV function	See detailed discussion of specific signs
Size/area	Cardiac size	See detailed discussion of specific signs
Character (thrusting, sustained, tapping, retracting)	A variety of abnormalities of LV structure and function	Single studies suggest potential utility in several contexts ²
Parasternal impulses	Abnormalities of cardiac and valvular function	Single studies suggest potential utility in several contexts ²
Palpable pulmonary second sound	Pulmonary hypertension	A single study suggests value in excluding PHT in mitral stenosis ⁵⁶
Auscultation:		
Intensity of first heart sound	Closure of mitral and tricuspid valves	
Loud/quiet/inaudible		Single studies suggest utility in several contexts ^{57,59}
Variable		A single study suggests utility in tachycardia diagnosis ³¹
Splitting of first heart sound	Bundle branch block, extrasystoles	No documented utility
Intensity of second heart sound	Closure of pulmonary and aortic valves	
Loud pulmonary second sound	Pulmonary hypertension	No discriminatory value for detection of PHT ⁶⁰
Quiet/inaudible aortic second sound	Abnormal closure of the aortic valve	See detailed discussion of specific signs
Splitting of second heart sound (physiological/fixed/reversed)	A variety of valvular and congenital abnormalities	Absence of fixed splitting useful in excluding atrial septal defect ⁶¹
Third heart sound	Increased early diastolic filling	See detailed discussion of specific signs

Table 1 | continued

Sign or manoeuvre	What is this sign purported to indicate?	Suggested clinical value
Fourth heart sound	Late diastolic filling related to atrial systole	Low specificity and sensitivity
Systolic/ejection clicks	A variety of valvular abnormalities	See detailed discussion of specific signs
Diastolic/opening snap	Pliant mitral valve leaflet in mitral stenosis	No data on diagnostic value. Proximity to second sound correlates with severity of mitral stenosis
Systolic murmur	A variety of abnormalities of central CVS blood flow	
Ejection/pan/late		See detailed discussion of specific signs
Pattern of radiation		See detailed discussion of specific signs
Effect of respiration	Increases or decreases venous return	
Louder on inspiration	Suggests murmur origin in right heart	LR+ of <5 in two studies of detection of tricuspid regurgitation and pulmonary stenosis ²
Effect of other manoeuvres (Valsalva/posture change/hand grip/leg elevation/amyl nitrite)	Increase or decrease venous return or SVR	LR+ of >5 in small number of studies in variety of diagnostic contexts ²
Diastolic murmur	A variety of abnormalities of central CVS blood flow	See detailed discussion of specific signs
Decrescendo/early	Aortic or pulmonary regurgitation	
Mid	Mitral or tricuspid stenosis	
Pericardial rub	Pericardial inflammation	The finding is pathognomic of pericarditis. As such it is of high specificity, but sensitivity is unknown
Peripheries and other organ systems		
Xanthelasmata	Dyslipidaemia	Presence independently predicts increased risk of IHD and MI ⁶²
Arcus lipidus	Dyslipidaemia	Of no independent diagnostic or prognostic value ⁶²
Transverse earlobe crease		Presence independently predicts increased risk of IHD and MI ⁶²
Male pattern baldness		Presence independently predicts increased risk of IHD and MI ⁶²
Hair greying		Of no independent diagnostic or prognostic value ⁶²
Cutaneous features of arterial insufficiency		See detailed discussion of specific signs
Oedema	Extravascular fluid overload	See detailed discussion of specific signs
Pulmonary rales (crepitations)	Interstitial fluid	Low sensitivity and specificity in diagnosis of CHF in isolation
Hepatomegaly		Low sensitivity and specificity for CHF. When pulsatile, strongly supports diagnosis of tricuspid regurgitation ²⁹
Splenomegaly		May provide supportive evidence of infective endocarditis, but non-specific and of low sensitivity
Finger clubbing		May provide supportive evidence of infective endocarditis, but non-specific and of low sensitivity. Of moderate sensitivity and specificity in some forms of cyanotic heart disease
Funduscopy		Roth spots may provide supportive evidence of infective endocarditis, but non-specific and of low sensitivity. Hypertensive retinopathy provides evidence of target organ damage and some features predict increased risk of CVS disease ⁶³

AR=aortic regurgitation; AS=aortic stenosis; CHF=congestive heart failure; CVS=cardiovascular system; HCM=hypertrophic obstructive cardiomyopathy; IHD=ischaemic heart disease; LR=likelihood ratio; LV=left ventricular; MI=myocardial infarction; PAD=peripheral arterial disease; PHT=pulmonary hypertension; SVR=systemic vascular resistance.

criteria for more detailed consideration are discussed and referenced in detail in the next section and indicated as such in the table.

Detailed discussion of selected signs

The signs included for detailed discussion in this section meet each of the following criteria:

- More than one study exists from which conventional indices of diagnostic accuracy can be derived.
- Where available, the positive likelihood ratio (LR+) for the sign is 5 or more, the negative likelihood ratio (LR-) is less than 0.2, or both. These values categorise a test as providing strong diagnostic evidence,⁶⁴ and they increase or decrease pre-test probability by around 30% (fig 5).
- The information obtained is of practical assistance in patient care, taking into account the additional information that is available or can be accessed through history, bedside tests, and additional investigations.

We stress that the exclusion of a sign from this section does not imply that the sign is of no value but simply that there is less evidence of its value than for signs that are included. For example, a pericardial rub is pathognomic of pericarditis, but no studies quantify its diagnostic value.

Jugular venous pulse and estimation of central venous pressure

Table 2 summarises studies evaluating the value of examination of the jugular venous pulse (JVP) as a means of assessing the central venous pressure (CVP).

The difficulties in collating studies of clinical assessment of the JVP have been described.⁷⁴ Differences in the preferred anatomical reference point, side of the neck used, preference for internal or external jugular vein, position of the patient, experience of observers, and clinical status of studied patients, including their volume status, severity of illness, presence of percutaneous cannulas in the veins assessed, and whether patients were breathing spontaneously or with mechanical support, all pose specific problems. In a minority of patients in these studies and in clinical practice, no attempt at pressure measurement can be made as the veins cannot be visualised (for example, in obese patients), but this proportion may decline with observers' training and experience.

Several studies suggest that clinical measurement of the JVP tends to underestimate the CVP measured by cardiac catheterisation. In some studies, this may reflect underestimation of the distance between the classical anatomical reference point of the sternal angle and the central right atrium, now recognised as 8 cm rather than

Table 2 | Studies of clinical assessment of jugular venous pressure (JVP) for estimation of central venous pressure (CVP)

Reference	What was done	Patients	Clinicians	No of patients (observations)	Summary of study conclusions
Davison, ⁶⁵ 1974, USA	External and internal JVP versus invasive CVP	“Seriously ill”	Two clinicians	39 (128)	IJV not usually visible. EJV measurement correlated poorly with CVP
Connors, ⁶⁶ 1983, USA	JVP versus invasive CVP	ICU	Medical student to attending physician	62 (310)	JVP underestimated CVP in ICU patients
Ducas, ⁶⁷ 1983, USA	JVP and HJR versus invasive CVP	Not defined	Not defined	48	CVP predicted from JVP in 44/48 comparisons
Cook, ⁶⁸ 1990, USA	JVP versus invasive CVP	ICU	15 different clinicians; medical student to staff physician	50 (150)	JVP inaccurate in ICU patients
Stein, ⁶⁹ 1997, USA	JVP versus IVC ultrasound RAP and invasive CVP	Patients with severe CHF	Not stated	22	Clinical estimates of RAP from JVP were accurate when RAP was normal, but systematically underestimated elevated RAP
Vinayak, ⁷⁰ 2006, USA	External JVP versus invasive CVP	Critically ill in ICU	Student to attending	35 (118)	JVP using EJV highly reliable for assessing elevated CVP (AUC 0.97, 95% CI 0.92 to 1.00) by attending physicians
Sinialo, ⁷¹ 2007, Finland	JVP with patient sitting versus invasive CVP	Pre-cardiac catheterisation	Two cardiologists	96	JVP using IJV in sitting patients had sensitivity of 65% and specificity of 85% to identify elevated CVP
Brennan, ⁷² 2007, USA	JVP versus invasive CVP and indirect comparison with ultrasound IVC	Pre-cardiac catheterisation	Four internal medicine residents	40 (44)	IJV not identified in 37% of patients. Sensitivity of JVP for elevated CVP 14%
Deol, ⁷³ 2011, USA	External and internal JVP clinically and by ultrasound versus invasive CVP	ICU	Not defined	38	EJV better than IJV in correlation with CVP, but ultrasound superior to both. Clinical and ultrasound measurement both underestimated CVP.
Rizkallah, ³⁰ 2014, Canada	JVP clinically versus ultrasound RAP	Routine echocardiography patient population	One student, one resident, one fellow	325	JVP sensitivity 86% by cardiology fellow

AUC=area under curve; CHF=congestive heart failure; EJV=external jugular vein; HJR=hepatojugular reflux; ICU=intensive care unit; IJV=internal jugular vein; IVC=inferior vena cava; RAP=right atrial pressure.

5 cm.⁷⁵ Although the right internal jugular vein has traditionally been regarded as the most accurate manometer, studies using either the left internal jugular vein or either external jugular vein have yielded accurate results.⁷⁶

On the basis of the available literature, determining whether the JVP, and by inference the CVP, is high, normal, or low, is preferable to attempting to define a specific value of the CVP derived from the JVP measurement. If the clinical measurement is made from the sternal angle, in whatever position the top of the pulsating venous column can be seen, a JVP of more than 3 cm above the sternal angle has an LR+ of 10.4 (95% confidence interval 5.5 to 19.9) and an LR- of 0.1 (0 to 0.6) for a CVP of more than 12 cm H₂O.² In a single study, in which the external jugular vein was used, a JVP of less than 3 cm above the sternal angle had an LR+ of 8.4 (2.8 to 25) and an LR- of 0.1 (0 to 0.7) for a CVP below 5 cm H₂O.⁷⁰ The abdominojugular test (hepatojugular reflux) may also be performed when examining the jugular veins and has an LR+ of 8.0 (no confidence interval) for elevated left heart filling pressures.⁷⁷⁻⁷⁹ The JVP is discussed further in the section on heart failure.

Signs of peripheral perfusion and shock syndromes

The diagnostic accuracy of physical examination markers of poor peripheral perfusion has been assessed in comparison with invasive haemodynamic indices in an intensive care setting. In a sub-study of 513 patients in intensive care with acute lung injury randomised to receive a pulmonary artery catheter, the combined presence of three physical examination findings (capillary refill time >2 s, knee mottling, or cool extremities) was highly specific but insensitive for identifying low cardiac output (<2.5 L/min/m²; specificity 98%; sensitivity 12%; LR+ 7.5) and low mixed venous oxygen saturation (<60%; specificity 99%; sensitivity 8%; LR+ 8.0).⁸⁰

In intensive care patients with septic shock, capillary refill time greater than 2.4 s at the fingertip (LR+ 3.0;

LR- 0.25) or less than 4.9 s at the knee (LR+ 5.1; LR- 0.21) six hours after diagnosis was predictive of death at 14 days.⁸¹ In a large study of consecutive acute adult medical admissions, prolonged capillary refill time on admission predicted one day and seven day mortality, independently of other demographic and clinical variables, with odds ratios of 1.69 (95% confidence interval 1.20 to 2.39; P=0.003) and 1.38 (1.12 to 1.69; P=0.002), respectively.⁸²

Physical examination signs of impaired peripheral perfusion may therefore hold both diagnostic and prognostic value in shock syndromes. Accordingly, assessment of capillary refill time and other markers of peripheral perfusion is recommended for the initial rapid evaluation of circulatory status in a wide variety of settings.^{83,84}

Pulsatile abdominal mass and abdominal aortic aneurysm

In studies in which aneurysmal dilatation is defined as greater than 3 cm, a pulsatile abdominal mass has an LR+ of between 8.0 (4.2 to 15.3)⁸⁵⁻⁹⁰ and 12 (7.4 to 19)⁹¹ for the presence of an aneurysm of this dimension. In general, the higher the aortic diameter used to define aneurysmal dilatation and the lower the abdominal girth,⁹² the higher the sensitivity of palpation. Therefore, a finding of a pulsatile mass in the abdomen on opportunistic physical examination suggests the presence of an abdominal aortic aneurysm and should prompt consideration of imaging.⁹³

Arterial pulse rhythm and atrial fibrillation

Studies of the value of pulse palpation as a screening tool for atrial fibrillation are comparatively large and have been included in systematic reviews and consensus guidance.^{94,95} Data pooled from three major studies indicate that the opportunistic finding of “any pulse irregularity” has an LR+ of 3.3 (3.0 to 3.7) for the presence of atrial fibrillation on electrocardiography.⁹⁶⁻⁹⁸ When such

irregularity is absent, this has an LR⁻ of 0.1 (0.1 to 0.2). This modest positive diagnostic value is greatly increased if the definition used is “continuous pulse irregularity,” when the LR⁺, derived from one large study,⁹⁶ rises to 24.1 (15.2 to 38). On the basis of these findings, pulse palpation for detection of atrial fibrillation is likely to have the greatest practical utility when used as an initial opportunistic screening method, identifying patients who need an electrocardiogram to confirm the underlying rhythm.

Absent peripheral pulses and bruits and peripheral arterial disease

A systematic review and subsequent studies have investigated the value of peripheral pulse palpation and auscultation in detecting peripheral arterial disease (PAD).²⁸⁻¹⁰⁰ In patients with symptoms suggestive of PAD, such as claudication, a palpable pulse abnormality and a lower limb bruit both have significance (LR⁺ 4.70 (2.20 to 9.90) and 5.60 (4.70 to 6.70), respectively).⁹⁹ Cool skin (LR⁺ 5.90, 4.10 to 8.60) and wounds or sores on the foot (LR⁺ 5.90, 2.60 to 13.40) also support the diagnosis, but other features such as hair loss, skin colour, and Buerger’s test are of little or no value.⁵⁰⁻¹⁰¹ In asymptomatic people, a femoral bruit has an LR⁺ of 4.80 (2.40 to 9.50) and “any pulse abnormality” has an LR⁺ of 3.10 (1.40 to 6.60). The absence of these signs does not exclude PAD but reduces the likelihood of severe disease.⁹⁹

Thus, the presence of abnormal physical examination findings may help to confirm the need for further investigation in patients with symptoms compatible with PAD. Insufficient evidence exists to support the use of physical examination in isolation as a screening tool in asymptomatic people, but further investigation may be reasonable following opportunistic detection of abnormalities in patients at moderate to high risk of PAD.

Pulse deficits and aortic dissection

A palpable pulse deficit of the carotid or upper limb pulses in patients with clinical presentations suggestive of acute thoracic aortic dissection supports the diagnosis, with a pooled LR⁺ of 6.0 (no confidence interval presented).²⁻¹⁰³ Patients with aortic dissection who have pulse deficits have more in-hospital complications and have higher mortality.¹⁰⁴ Aortic dissection is underdiagnosed, and the absence of these findings should never dissuade a clinician from the diagnosis. All clinical clues may be of value, and relevant imaging should always be undertaken when clinical suspicion is present.

Abdominal bruits and renovascular hypertension

In patients with hypertension, abdominal bruits with systolic-diastolic components are strongly supportive of a diagnosis of renovascular disease (LR⁺ 39; LR⁻ 0.6; no confidence interval presented)¹⁰⁵ and bruits with isolated systolic components less so (LR⁺ 5.6, 4 to 7.7; LR⁻ 0.6, no confidence interval presented).¹⁰⁶⁻¹⁰⁹ Accordingly, auscultation for abdominal bruits may help to identify patients who need renal Doppler ultrasound. However, the absence of bruits does not exclude the presence of renal artery stenosis.

Carotid bruits and carotid stenosis

Meta-analysis suggests that carotid bruits are of moderate value in the detection of clinically relevant (>70%) carotid stenosis, with pooled sensitivity of 0.53 (0.50 to 0.55), specificity of 0.83 (0.82 to 0.84), and diagnostic odds ratio of 4.32 (2.78 to 6.66).¹¹⁰ Patients with known or suspected carotid territory cerebrovascular disease should undergo appropriate imaging of the vessel regardless of carotid auscultatory findings, as absence of a bruit does not exclude disease. However, opportunistic detection of a bruit in other settings should prompt consideration of further investigation, as detection of significant carotid stenosis may have important therapeutic implications such as the need for secondary preventive measures.¹¹¹

Physical examination in heart failure

Cardiac failure is associated with several characteristic physical examination findings. Salt and water retention, together with elevated ventricular filling pressures, may lead to signs of congestion such as an elevated JVP, peripheral (usually lower limb) oedema, pulmonary rales (crepitations), and, less frequently, ascites or hepatomegaly. A third heart sound caused by abrupt deceleration of rapid ventricular (usually left ventricular) filling occurs when the rate of filling exceeds ventricular compliance and usually signifies the combination of elevated mean left atrial pressure and left ventricular dysfunction. A displaced apex beat suggests structural left ventricular remodelling and correlates with increased left ventricular end diastolic volume and mass, as well as reduced left ventricular ejection fraction.

Table 3 summarises the clinical value of the physical examination of the cardiovascular system in patients with known or suspected heart failure. One difficulty in evaluating the diagnostic value of physical examination findings in cardiac failure lies in the lack of a single diagnostic test to use as a reference standard. Guidelines recommend diagnosis based on a combination of typical symptoms, physical examination findings (described above), and demonstration of a relevant abnormality of cardiac structure and/or function.¹²⁴⁻¹²⁵ For the studies of diagnostic accuracy in table 3, the reference standard for “heart failure” was therefore final consensus diagnosis based on a combination of all available clinical and investigational data. Studies using left ventricular systolic dysfunction (LVSD) as a reference standard may tend to underestimate the diagnostic value of physical examination findings, as the clinical syndrome of heart failure may occur in the absence of LVSD. However, we have presented data for the diagnostic accuracy of physical examination findings for detecting LVSD in a primary care setting.

In general, physical examination findings in heart failure lack sensitivity, and their absence has minimal value in excluding the diagnosis (table 3). Additionally, dependent oedema and pulmonary rales are relatively non-specific and have, at best, modest value for ruling in heart failure. However, elevated JVP, third heart sound, and displacement of the apex beat are each highly specific and substantially increase the likelihood of heart failure (table 3). When present, a third heart

STATE OF THE ART REVIEW

Table 3 | Clinical value of physical examination of the cardiovascular system in patients with known or suspected heart failure (HF)

Clinical context	Raised JVP	Peripheral oedema	Rales	Third heart sound	Displaced apex	Comments
Diagnostic accuracy						
Pooled LR+ (CI) for HF in emergency department patients with dyspnoea ¹¹²	5.1 (3.2 to 7.9)	2.3 (1.5 to 3.7)	2.8 (1.9 to 4.1)	11.0 (4.9 to 25)		LR- >0.5 for all except displaced apex
Pooled LR+ (CI/range) for LVSD in community patients with suspected HF ¹¹³	4.36 (range 2.7-7.4)	1.18 (range 1.0-1.5)	1.53 (1.2 to 2.2)	7.34 (range 1.6-32.4)	16.0 (8.2 to 30.9)	LR- >0.5 for all
LR+ range for HF (any ejection fraction) in community patients with suspected HF	4.8 to 7.5 ^{114,115}	1.9 to 3.0 ^{114,115}	3.0 to 3.4 ^{114,115}	22.3 ¹¹⁵	6.7 ¹¹⁵	LR- >0.5 for all
Independent prognostic value						
Asymptomatic LVSD ^{†116} :						
Progression to clinical HF	1.51 (1.04 to 2.19)			1.38 (1.09 to 1.73)		Patients with third heart sound also at increased risk of hospital admission for HF: 1.47 (1.06 to 2.05)
Death or development of HF	1.54 (1.11 to 2.12)			1.34 (1.09 to 1.64)		
Suspected HF in community:						
All cause mortality ^{‡114}	1.45 (1.16 to 1.83) for rales plus JVP or oedema					Congestion stronger predictor of outcomes than LVEF. Multivariate analysis included NT-proBNP
Stable CHF (NYHA I-IV) ^{†117} :						
Hospital admission for HF	1.32 (1.08 to 1.62)			1.42 (1.21 to 1.66)		All cause death: 1.17 (1.02 to 1.35) for JVP and third heart sound
Death or hospital admission for HF	1.30 (1.11 to 1.53)			1.40 (1.14 to 1.71)		
Stable CHF (NYHA III-IV) ^{†118} :						
All cause mortality	1.52 (1.22 to 1.91)	NS				1.61 (1.18 to 2.20) for oedema + JVP
Hospital admission for HF	1.21 (1.02 to 1.43)	1.25 (1.06 to 1.48)				1.37 (1.08 to 1.73) for oedema + JVP
Stable CHF with AF ^{†119} :						
CV mortality	NS	1.25 (1.0 to 1.57)	1.41 (1.07 to 1.86)	NS		JVP, oedema, rales, and third heart sound each strongly predictive of all cause death on univariate analysis
Hospital admission for HF	NS	NS	1.42 (1.05 to 1.90)	NS		
Acute decompensated HF:						
In-hospital mortality ^{‡120}	NS	NS	NS	1.69 (1.19 to 2.41)		JVP and oedema form part of composite congestion scores that predict post-discharge mortality
Post-discharge mortality ^{‡121,123}	See comment	See comment				

CHF=congestive heart failure; JVP=jugular venous pressure; LR=likelihood ratio; LVEF=left ventricular ejection fraction; LVSD=left ventricular systolic dysfunction; NT-proBNP=N-terminal pro-hormone of brain natriuretic peptide; NYHA=New York Heart Association.

*All results based on multivariate analysis including clinical, biochemical, electrocardiographic, and echocardiographic variables +/- NT-proBNP. Data expressed as hazard ratios unless otherwise stated.

†Relative risk (95% CI).

‡Odds ratio (95% CI).

sound strongly supports the diagnosis of heart failure, although it is detected infrequently in community settings (1-5%), which may limit its practical value. In contrast, a displaced apex beat was the most helpful physical examination finding in a meta-analysis of primary care studies and, in a subsequent study, retained substantial diagnostic value after adjustment for other physical examination findings and symptoms (odds ratio 5.4, 2.8 to 10.5). A raised JVP increases the probability of heart failure across all clinical settings and, in primary care, has diagnostic value independent of all other clinical variables and electrocardiography findings (odds ratio 8.7, 2.6 to 29.2).¹²⁶

In patients with stable or decompensated heart failure, signs of congestion or the presence of a third heart sound predict a range of adverse outcomes independently of other established prognostic variables including age, New York Heart Association class, and left ventricular ejection fraction (table 3). Relief of congestion is a key therapeutic goal in patients with heart failure and requires titration of therapies to physical examination findings.^{124,125} Importantly, effective relief of congestion (assessed using composite scores incorporating JVP and/or oedema) in patients admitted to hospital reduces the risk of subsequent adverse outcomes compared with those with residual congestion.^{113,114}

Abnormal systolic murmur and valvular heart disease

As systolic murmurs are often detected in patients without significant valvular or other structural heart disease,¹²⁷ an important aspect of clinical evaluation is to

distinguish such “functional” murmurs from pathological murmurs. Guidelines do not recommend echocardiographic evaluation of suspected functional murmurs in asymptomatic people.^{128,129} In four separate studies of patients with a systolic murmur examined by a cardiologist, classification of the murmur as “abnormal” significantly increased the likelihood of valve disease, with likelihood ratios ranging from 3.8 to infinity, whereas classification as “normal” reduced the likelihood (likelihood ratios of 0-0.3).¹³⁰⁻¹³³

The physical examination, whether performed by cardiologists or non-cardiologists, is also useful for diagnosing valvular heart disease in asymptomatic patients without known cardiac disease and in patients attending the emergency department,^{134,135} with a pooled LR+ of 15 (11 to 20)³ for an “abnormal” murmur and 0.25 (0.17 to 0.36) for a functional murmur or no murmur. Substantial agreement existed between non-cardiologist emergency department physicians for identification of a systolic murmur ($\kappa=0.8$).¹²⁷

Intensity and duration of the murmur are the most helpful features to differentiate functional from pathological murmurs. A loud murmur and holosystolic timing help to rule in and rule out underlying valve disease, with an LR+ of 6.5 (2.3 to 19) and 8.7 (2.3 to 33) and an LR- of 0.08 (0.02 to 0.31) and 0.19 (0.08 to 0.43), respectively.¹³⁰ A systolic thrill is strongly suggestive of significant valve disease, but its absence is not useful in ruling out a pathological murmur (LR+ 12, 0.76 to 205; LR- 0.73, 0.58 to 0.93).¹³⁰

Table 4 | Clinical features that influence the likelihood of aortic stenosis in patients with a systolic murmur

Clinical feature	Significance (95% CI)
Findings that make moderate to severe aortic stenosis less likely	
Murmur not heard over right clavicle	LR– 0.10 (0.02 to 0.44) ¹³⁷
Lack of carotid radiation	LR– 0.1–0.29 ²⁹⁻¹³⁷
Mid-systolic timing	LR– 0.05 (0 to 0.3) ²⁹
Grade 1 intensity	LR– 0.2 (0.1 to 0.50) ²⁹
Findings that make moderate to severe aortic stenosis more likely	
Delayed upstroke of carotid pulse	(LR+ 6.8 (4 to 11.5) to 130 (33 to 560)) ²⁹⁻¹³⁷
Reduced or absent second heart sound	(LR+ 7.5 (3.2 to 17) to 50 (24 to 100)) ²⁹⁻¹³⁷
Carotid radiation	(LR+ 8.1 (4 to 16) to 12.4 (4.5 to 34)) ²⁹⁻¹³⁷
Reduced carotid volume	(LR+ 2.0 (1 to 3.2) to 2.3 (1.7 to 3.0)) ²⁹⁻¹³⁷

LR–=negative likelihood ratio; LR+=positive likelihood ratio.

Auscultation for a systolic murmur is therefore helpful in determining the likelihood of valvular heart disease and allows patients with suspected valve disease to receive further investigation (usually echocardiography) to confirm the nature and severity of valve pathology while avoiding unnecessary investigation in patients with low likelihood of valve disease.

Characteristic murmurs and specific valve lesions

Aortic stenosis

In most studies, the absence of a systolic murmur effectively excludes the diagnosis of significant aortic stenosis, with negative likelihood ratios ranging from 0 to 0.05.²⁹⁻¹³⁷ In a cohort of patients admitted with hip fracture, the LR– for moderate to severe aortic stenosis when no murmur was heard was 0.36 (0.20 to 0.62). However, the negative predictive value remained high (96.8% for moderate or greater stenosis; 99.6% for severe stenosis).¹³⁸ Among patients with a systolic murmur, several clinical signs have diagnostic value for aortic stenosis (table 4).

None of these, in isolation, has sufficient accuracy to rule in aortic stenosis, but moderate to severe stenosis is highly likely if three or more are present (LR+ 40, 6.6 to 239).¹³⁷ Among non-cardiologist examiners, moderate agreement existed for detection of a soft or absent second heart sound ($\kappa=0.54$) and fair agreement for identification of radiation to right clavicle ($\kappa=0.36$), radiation to right carotid ($\kappa=0.33$), delayed carotid upstroke ($\kappa=0.26$), and reduced carotid volume ($\kappa=0.24$).¹³⁷

Mitral regurgitation

In unselected hospital inpatients undergoing echocardiography, a murmur that extends from the apex to at least the anterior axillary line suggests moderate or greater mitral regurgitation (LR+ 6.8, 3.9 to 11.9).²⁹ Importantly, the absence of a murmur is not helpful in excluding mitral regurgitation in the setting of acute myocardial infarction (LR– 0.66, 0.25 to 1.0).¹³⁹ However, in patients with known mitral valve prolapse, the absence of a holosystolic or late systolic murmur, identified by a cardiologist, makes moderate or severe mitral regurgitation unlikely (LR– 0, 0 to 0.8).¹⁴⁰ The intensity of the murmur correlates with the severity of regurgitation. Very loud murmurs (grade 4–5) substantially increase the likelihood of severe mitral regurgitation (LR+ 14, 3.3 to 56), whereas grade 1–2 murmurs are unlikely to reflect severe regurgitation (LR+ 0.19, 0.11 to 0.33).²⁹

Tricuspid regurgitation

In patients referred for echocardiography, the presence of a characteristic murmur, as determined by a cardiologist, strongly predicts the presence of tricuspid regurgitation (LR+ 10.1, 5.8 to 18).¹⁴¹ However, the absence of a murmur is not helpful in excluding tricuspid regurgitation (LR– range 0.41–0.60).^{133 142}

Specific features of the murmur with diagnostic value include localisation to the lower left sternal edge (not audible above the third intercostal space or lateral to the mid-clavicular line), which strongly supports the diagnosis¹⁴² (LR+ 8.4, 3.5 to 20.3) and an increase in intensity with inspiration, which, among experienced cardiologists, distinguishes tricuspid from mitral regurgitation (LR+ ∞ , 3.1 to ∞ ; LR– 0.20, 0.07 to 0.45).¹⁴² Additionally, the presence of a giant V wave in the jugular veins helps to rule in the diagnosis of tricuspid regurgitation (LR+ 10.9, 5.5 to 22).¹⁴²

Aortic regurgitation

A systematic review indicates that auscultation for the typical early diastolic, high pitched decrescendo murmur of aortic regurgitation is helpful for both ruling in and ruling out aortic regurgitation.¹⁴³ In the two highest quality studies, the presence of a typical aortic regurgitation murmur had an LR+ of 8.8–32 for detecting aortic regurgitation of at least mild severity and 4.0–8.3 for moderate to severe aortic regurgitation.^{144 145} Conversely, the absence of a typical aortic regurgitation murmur had an LR – of 0.2–0.3 for excluding any aortic regurgitation and 0.1 (0 to 0.3) for moderate to severe aortic regurgitation. The clinical context of a diastolic murmur is important, as patients with end stage renal failure and volume overload may have transient early diastolic murmurs that resolve with correction of volume overload.¹⁴⁶

The physical examination is of limited value in assessing the severity of aortic regurgitation. However, severe aortic regurgitation is unlikely with very quiet (grade 1) murmurs (LR+ 0, 0 to 0.9) and more likely when either a third heart sound (LR+ 5.9, 1.4 to 25) or grade 3 murmur (LR+ 4.5, 1.6 to 14) is present.^{44 147}

Overall, these findings indicate that physical examination findings are valuable in identifying or excluding the presence of specific valve lesions. Although most patients with suspected valve disease in contemporary practice should undergo formal evaluation with echocardiography, identification of the likely underlying lesion may help to inform the urgency of echocardiography or guide initial management when echocardiography is not immediately available. Specific clinical settings in which insight from the physical examination may be particularly helpful include patients needing urgent surgery, those with suspected endocarditis, and those with new onset of decompensated heart failure.

Clinical context

Physical examination is immediately available, rapid and repeatable, relatively inexpensive, safe, and non-invasive. The data presented in this review provide evidence that some of its elements can still support the investigation and ongoing care of patients with known or suspected cardiovascular disease in contemporary healthcare.

GLOSSARY OF ABBREVIATIONS

CVP	central venous pressure
JVP	jugular venous pressure
LR–	negative likelihood ratio
LR+	positive likelihood ratio
LVSD	left ventricular systolic dysfunction
PAD	peripheral arterial disease
TAD	technological aids to diagnosis

In emergency care, physical examination of the cardiovascular system can direct initial treatment in the period before TAD can be accessed, determine the choice and urgency of investigation, and support monitoring of response to treatment. A firm clinical impression of decompensated heart failure in the emergency department does not obviate the need for echocardiography but does permit prompt starting of diuretic and vasodilator therapy, as well as avoidance of potentially inappropriate treatments. Once a diagnosis is established, repeated physical examination can help in optimising management, most notably through identifying the presence of residual congestion that, if untreated, is associated with repeated hospital admission and poorer survival.

In the community setting, or other settings where TAD are less readily available, the physical examination holds particular value in rationalising and targeting specialist investigation. An abnormal murmur in a patient with dyspnoea establishes the need for echocardiography, and the coexistence of physical findings of heart failure would mandate urgent referral. The physical examination also provides a simple method for opportunistic detection of asymptomatic but potentially important disease such as atrial fibrillation, valvular heart disease, and abdominal aortic aneurysm, which can then be assessed further by appropriate investigation.

Conclusions

Despite its apparent benefits, physical examination is increasingly poorly practised, understood, and trusted in some contemporary healthcare settings. These changes have occurred despite continued focus on the teaching and assessment of relevant skills in undergraduate medical education, although this is not always sustained into postgraduate education with similar intensity.¹⁴⁸ The use of TAD has increased substantially over the same period that the practice of the physical examination of the cardiovascular system has declined. However, this may reflect factors not primarily related to the diagnostic accuracy of physical examination, such as models of reimbursement of physicians and patients' expectations, rather than considered comparisons of the efficacy of diagnostic strategies that use physical examination and TAD alone or in combination.¹⁴⁹⁻¹⁵¹

Whatever the explanations for the relative change in use, concerns have arisen about the implications. Underuse or absence of physical examination can lead to misdiagnosis and may adversely affect the doctor-patient relationship.^{23,152} Overuse of TAD can lead to unwarranted financial expense and physical harm to patients. Major professional organisations now stress the potential role of physical examination in reducing unnecessary technological testing,¹²⁸ and initiatives such as Choosing Wisely promote more judicious selection of investigations (<http://www.choosingwisely.org>).

RESEARCH QUESTIONS

- Cost effectiveness of strategies that combine physical examination and technological aids to diagnosis (TAD) versus those using TAD alone
- Relation between use of physical examination and use of TAD
- Patients' attitudes to care strategies including and excluding physical examination
- Diagnostic error rate in care strategies that include and exclude physical examination
- Diagnostic value of combinations of signs or of signs and symptoms
- Value accorded to physical examination by physicians in contemporary practice
- Diagnostic accuracy of signs of cardiac disease now uncommon in developed countries

However, whether improved physical examination, which is not without its own financial cost, can effectively reduce diagnostic error, the overuse of TAD, or the costs of care is not known. Although enhanced physical examination skills could lead to improved confidence in bedside decision making, clinicians' understanding of statistical data derived from clinical assessment is often poor,⁴⁰ and intolerance of perceived "uncertainty" in the absence of the results of a TAD may still drive use, even in situations in which subsequent diagnostic probability may be unaffected.¹⁵³ Whether patients will now trust judgments based on clinical assessment alone or be comfortable with treatment decisions made in the absence of TAD is also uncertain.

Furthermore, the increasing availability and decreasing cost of point of care TAD, such as portable bedside handheld ultrasound, is likely to further question the value of traditional physical examination techniques, particularly in the elucidation and detection of murmurs and evaluation of CVP.¹⁵⁴ However, the costs of providing such technology to clinicians, delivering the education and training required to implement and sustain widespread use, and the diagnostic accuracy of bedside ultrasound during routine clinical care in non-specialist settings have yet to be defined or compared with those of the physical examination. Such background should provide a continuing incentive to reappraise the value of the physical examination.

Given the infrequent occurrence of some signs, the difficulty in ensuring that observers' ability and technique are standardised, and the necessity to include large numbers of observations to generate robust estimates of diagnostic accuracy, it is unlikely that further research based around the simple association of specific signs with specific diagnoses is likely to be practicable or informative. Future research may thus be more usefully focused on the areas listed in the "Research questions" box.

We could not have completed this review without the existing published work and helpful comments of Steven McGee (Seattle). We thank Stuart Shaw (Edinburgh) and Junaid Zaman (Stanford) for their comments and Estela Dukan (Edinburgh) for helping with the literature search.

Contributors: AE, AJ, and AV all contributed to the creation of the outline and drafts of the paper. AE and AJ led the literature review. AE is the guarantor.

Competing interests: We have read and understood the BMJ policy on declaration of interests and declare the following interests: none.

Provenance and peer review: Commissioned; externally peer reviewed.

Patient involvement: No patients were asked for input in the creation of this article.

- 1 Aird WC. Discovery of the cardiovascular system: from Galen to William Harvey. *J Thromb Haemost* 2011;9(Suppl 1):118-29. doi:10.1111/j.1538-7836.2011.04312.x pmid:21781247.
- 2 McGee S. *Evidence-based Physical Diagnosis*. 3rd ed. WB Saunders, 2012;doi:10.1016/B978-1-4377-2207-9.00001-X.
- 3 Simel DL, Rennie D, Keitz SA, eds. *The rational clinical examination: Evidence based clinical diagnosis*. McGraw-Hill Companies, 2008.
- 4 Roelandt JR. The decline of our physical examination skills: is echocardiography to blame? *Eur Heart J Cardiovasc Imaging* 2014;15:249-52. doi:10.1093/ehjci/jet195 pmid:24282219.
- 5 Tavel ME. Cardiac auscultation. A glorious past-but does it have a future? *Circulation* 1996;93:1250-3. doi:10.1161/01.CIR.93.6.1250 pmid:8653848.
- 6 Craige E. Should auscultation be rehabilitated? *N Engl J Med* 1988;318:1611-3. doi:10.1056/NEJM198806163182409 pmid:3374530.
- 7 Fred HL. Hyposkillia: deficiency of clinical skills. *Tex Heart Inst J* 2005;32:255-7. pmid:16392201.
- 8 Vukanovic-Criley JM, Criley S, Warde CM, et al. Competency in cardiac examination skills in medical students, trainees, physicians, and faculty: a multicenter study. *Arch Intern Med* 2006;166:610-6. doi:10.1001/archinte.166.6.610 pmid:16567598.
- 9 Johnson JE, Carpenter JL. Medical house staff performance in physical examination. *Arch Intern Med* 1986;146:937-41. doi:10.1001/archinte.1986.00360170163023 pmid:3963985.
- 10 Mangione S, Nieman LZ. Cardiac auscultatory skills of internal medicine and family practice trainees. A comparison of diagnostic proficiency. *JAMA* 1997;278:717-22. doi:10.1001/jama.1997.03550090041030 pmid:9286830.
- 11 St Clair EW, Oddone EZ, Waugh RA, Corey GR, Feussner JR. Assessing housestaff diagnostic skills using a cardiology patient simulator. *Ann Intern Med* 1992;117:751-6. doi:10.7326/0003-4819-117-9-751 pmid:1416578.
- 12 Sztajzel JM, Picard-Kossovsky M, Lerch R, Vuille C, Sarasin FP. Accuracy of cardiac auscultation in the era of Doppler-echocardiography: a comparison between cardiologists and internists. *Int J Cardiol* 2010;138:308-10. doi:10.1016/j.ijcard.2008.06.066 pmid:18762344.
- 13 Paley L, Zornitzki T, Cohen J, Friedman J, Kozak N, Schattner A. Utility of clinical examination in the diagnosis of emergency department patients admitted to the department of medicine of an academic hospital. *Arch Intern Med* 2011;171:1394-6. doi:10.1001/archinternmed.2011.340 pmid:21824956.
- 14 Wells PS, Anderson DR, Rodger M, et al. Derivation of a simple clinical model to categorize patients probability of pulmonary embolism: increasing the models utility with the SimpliRED D-dimer. *Thromb Haemost* 2000;83:416-20. pmid:10744147.
- 15 Killip T 3rd, Kimball JT. Treatment of myocardial infarction in a coronary care unit. A two year experience with 250 patients. *Am J Cardiol* 1967;20:457-64. doi:10.1016/0002-9149(67)90023-9 pmid:6059183.
- 16 Smith ME, Chiovaro JC, O'Neil M, et al. Early warning system scores for clinical deterioration in hospitalized patients: a systematic review. *Ann Am Thorac Soc* 2014;11:1454-65. doi:10.1513/AnnalsATS.201403-1020C pmid:25296111.
- 17 Wells PS, Hirsh J, Anderson DR, et al. Accuracy of clinical assessment of deep-vein thrombosis. *Lancet* 1995;345:1326-30. doi:10.1016/S0140-6736(95)92535-X pmid:7752753.
- 18 Prendergast BD. Diagnostic criteria and problems in infective endocarditis. *Heart* 2004;90:611-3. doi:10.1136/hrt.2003.029850 pmid:15145855.
- 19 Bergan JJ, Schmid-Schönbein GW, Smith PD, Nicolaidis AN, Boisseau MR, Eklof B. Chronic venous disease. *N Engl J Med* 2006;355:488-98. doi:10.1056/NEJMra055289 pmid:16885552.
- 20 McKibbin KA, Walker-Dilks CJ. Beyond ACP Journal Club: how to harness MEDLINE for diagnostic problems. *ACP J Club* 1994;121(Suppl 2):A10-2. pmid:8069494.
- 21 Post TW, ed. UpToDate. Waltham, MA
- 22 Simel DL, Rennie D, Bossuyt PM. The STARD statement for reporting diagnostic accuracy studies: application to the history and physical examination. *J Gen Intern Med* 2008;23:768-74. doi:10.1007/s11606-008-0583-3 pmid:18347878.
- 23 Verghese A, Charlton B, Kassirer JP, Ramsey M, Ioannidis JP. Inadequacies of physical examination as a cause of medical errors and adverse events: A collection of vignettes. *Am J Med* 2015;128:1322-4.e3. doi:10.1016/j.amjmed.2015.06.004 pmid:26144103.
- 24 Cheraghi-Sohi S, Hole AR, Mead N, et al. What patients want from primary care consultations: a discrete choice experiment to identify patients' priorities. *Ann Fam Med* 2008;6:107-15. doi:10.1370/afm.816 pmid:18332402.
- 25 Verghese A, Brady E, Kapur CC, Horwitz RI. The bedside evaluation: ritual and reason. *Ann Intern Med* 2011;155:550-3. doi:10.7326/0003-4819-155-8-201110180-00013 pmid:22007047.
- 26 Wiley B, Mohanty B. Handheld ultrasound and diagnosis of cardiovascular disease at the bedside. *J Am Coll Cardiol* 2014;64:229-30. doi:10.1016/j.jacc.2014.05.011 pmid:25011727.
- 27 Armstrong DW, Tobin C, Matangi MF. The accuracy of the physical examination for the detection of lower extremity peripheral arterial disease. *Can J Cardiol* 2010;26:e346-50. doi:10.1016/S0828-282X(10)70467-0 pmid:21165366.
- 28 Cournot M, Boccalon H, Cambou J-PP, et al. Accuracy of the screening physical examination to identify subclinical atherosclerosis and peripheral arterial disease in asymptomatic subjects. *J Vasc Surg* 2007;46:1215-21. doi:10.1016/j.jvs.2007.08.022 pmid:18154997.
- 29 McGee S. Etiology and diagnosis of systolic murmurs in adults. *Am J Med* 2010;123:913-921.e1. doi:10.1016/j.amjmed.2010.04.027 pmid:20920693.
- 30 Rizkallah J, Jack M, Saeed M, Shafer LA, Vo M, Tam J. Non-invasive bedside assessment of central venous pressure: scanning into the future. *PLoS One* 2014;9:e109215. doi:10.1371/journal.pone.0109215 pmid:25279995.
- 31 Garratt CJ, Griffith MJ, Young G, et al. Value of physical signs in the diagnosis of ventricular tachycardia. *Circulation* 1994;90:3103-7. doi:10.1161/01.CIR.90.6.3103 pmid:7994860.
- 32 Marcus GM, Vessey J, Jordan MV, et al. Relationship between accurate auscultation of a clinically useful third heart sound and level of experience. *Arch Intern Med* 2006;166:617-22. doi:10.1001/archinte.166.6.617 pmid:16567599.
- 33 Wood PW, Choy JB, Nanda NC, Becher H. Left ventricular ejection fraction and volumes: it depends on the imaging method. *Echocardiography* 2014;31:87-100. doi:10.1111/echo.12331 pmid:24786629.
- 34 Unzek S, Popovic ZB, Marwick TH. Diastolic Guidelines Concordance Investigators. Effect of recommendations on interobserver consistency of diastolic function evaluation. *JACC Cardiovasc Imaging* 2011;4:460-7. doi:10.1016/j.jcmg.2011.01.016 pmid:21565732.
- 35 Eusebi P. Diagnostic accuracy measures. *Cerebrovasc Dis* 2013;36:267-72. doi:10.1159/000353863 pmid:24135733.
- 36 Whiting PF, Davenport C, Jameson C, et al. How well do health professionals interpret diagnostic information? A systematic review. *BMJ Open* 2015;5:e008155. doi:10.1136/bmjopen-2015-008155 pmid:26220870.
- 37 McGee S. Simplifying likelihood ratios. *J Gen Intern Med* 2002;17:646-9. doi:10.1046/j.1525-1497.2002.10750.x pmid:12213147.
- 38 Wildes T, Anderson R. The adult screening physical examination: what physicians do. *WJM* 2004;103:60-5. pmid:15101470.
- 39 Haring CM, van der Meer JW, Postma CT. A core physical examination in internal medicine: what should students do and how about their supervisors? *Med Teach* 2013;35:e1472-7. doi:10.3109/0142159X.2013.778396 pmid:23570566.
- 40 Mallett S, Halligan S, Thompson M, Collins GS, Altman DG. Interpreting diagnostic accuracy studies for patient care. *BMJ* 2012;345:e3999. doi:10.1136/bmj.e3999 pmid:22750423.
- 41 Arnell TD, de Virgilio C, Chang L, Bongard F, Stabile BE. Admission factors can predict the need for ICU monitoring in gallstone pancreatitis. *Am Surg* 1996;62:815-9. pmid:8813162.
- 42 Zhang D, Shen X, Qi X. Resting heart rate and all-cause and cardiovascular mortality in the general population: a meta-analysis. *CMAJ* 2016;188:E53-63. doi:10.1503/cmaj.150535 pmid:26598376.
- 43 McGee S, Abernethy WB 3rd, Simel DL. The rational clinical examination. Is this patient hypovolemic? *JAMA* 1999;281:1022-9. doi:10.1001/jama.281.11.1022 pmid:10086438.
- 44 Desjardins VA, Enriquez-Sarano M, Tajik AJ, Bailey KR, Seward JB. Intensity of murmurs correlates with severity of valvular regurgitation. *Am J Med* 1996;100:149-56. doi:10.1016/S0002-9343(97)89452-1 pmid:8629648.
- 45 Roy CL, Minor MA, Brookhart MA, Choudhry NK. Does this patient with a pericardial effusion have cardiac tamponade? *JAMA* 2007;297:1810-8. doi:10.1001/jama.297.16.1810 pmid:17456823.
- 46 Curtiss EI, Reddy PS, Uretsky BF, Cecchetti AA. Pulsus paradoxus: definition and relation to the severity of cardiac tamponade. *Am Heart J* 1988;115:391-8. doi:10.1016/0002-8703(88)90487-5 pmid:3341174.
- 47 Surawicz B, Fisch C. Cardiac alternans: diverse mechanisms and clinical manifestations. *J Am Coll Cardiol* 1992;20:483-99. doi:10.1016/0735-1097(92)90122-4 pmid:1634690.
- 48 Ashrafian H. Pulsatile pseudo-proptosis, aortic regurgitation and 31 eponyms. *Int J Cardiol* 2006;107:421-3. doi:10.1016/j.ijcard.2005.01.060 pmid:16503268.
- 49 Deakin CD, Low JL. Accuracy of the advanced trauma life support guidelines for predicting systolic blood pressure using carotid, femoral, and radial pulses: observational study. *BMJ* 2000;321:673-4. doi:10.1136/bmj.321.7262.673 pmid:10987771.
- 50 Insall RL, Davies RJ, Prout WG. Significance of Buerger's test in the assessment of lower limb ischaemia. *J R Soc Med* 1989;82:729-31. pmid:2693712.
- 51 Fields WS, Lemak NA. Joint Study of extracranial arterial occlusion. VII. Subclavian steal--a review of 168 cases. *JAMA* 1972;222:1139-43. doi:10.1001/jama.1972.03210090019004 pmid:4678043.
- 52 Danford DA, Fletcher SE, Martin AB, Gumbiner CH. Accuracy of clinical diagnosis of left heart valvular or obstructive lesions in pediatric outpatients with heart murmur. *Am J Cardiol* 2002;89:878-84. doi:10.1016/S0002-9149(02)02208-7 pmid:11909582.

- 53 Schmidt DE, Shah PK. Accurate detection of elevated left ventricular filling pressure by a simplified bedside application of the Valsalva maneuver. *Am J Cardiol* 1993;71:462-5. doi:10.1016/0002-9149(93)90458-0 pmid:8430644.
- 54 Rizkallah J, Jack M, Saeed M, Shafer LA, Vo M, Tam J. Non-invasive bedside assessment of central venous pressure: scanning into the future. *PLoS One* 2014;9:e109215. doi:10.1371/journal.pone.0109215 pmid:25279995.
- 55 Heckerling PS, Wiener SL, Moses VK, Claudio J, Kushner MS, Hand R. Accuracy of precordial percussion in detecting cardiomegaly. *Am J Med* 1991;91:328-34. doi:10.1016/0002-9343(91)90149-R pmid:1835287.
- 56 Whitaker W. Clinical diagnosis of pulmonary hypertension in patients with mitral stenosis. *Q J Med* 1954;23:105-12. pmid:13134464.
- 57 Tei C, Shah PM, Cherian G, Wong M, Ormiston JA. The correlates of an abnormal first heart sound in mitral-valve-prolapse syndromes. *N Engl J Med* 1982;307:334-9. doi:10.1056/NEJM198208053070602 pmid:7088098.
- 58 Wood P. An appreciation of mitral stenosis. I. Clinical features. *Br Med J* 1954;1:1051-63. doi:10.1136/bmj.1.4870.1051 pmid:13149899.
- 59 Wood P. An appreciation of mitral stenosis. II. Investigations and results. *Br Med J* 1954;1:1113-24. doi:10.1136/bmj.1.4871.1113 pmid:13149911.
- 60 Fowler NO, Noble WJ, Giarratano SJ, Mannix EP. The clinical estimation of pulmonary hypertension accompanying mitral stenosis. *Am Heart J* 1955;49:237-49. doi:10.1016/0002-8703(55)90196-4 pmid:13228356.
- 61 Perloff JK, Harvey WP. Mechanisms of fixed splitting of the second heart sound. *Circulation* 1958;18:998-1009. doi:10.1161/01.CIR.18.5.998 pmid:13585583.
- 62 Christoffersen M, Frikke-Schmidt R, Schnohr P, Jensen GB, Nordestgaard BG, Tybjaerg-Hansen A. Visible age-related signs and risk of ischemic heart disease in the general population: a prospective cohort study. *Circulation* 2014;129:990-8. doi:10.1161/CIRCULATIONAHA.113.001696 pmid:24334176.
- 63 Grosso A, Veglio F, Porta M, Grignolo FM, Wong TY. Hypertensive retinopathy revisited: some answers, more questions. *Br J Ophthalmol* 2005;89:1646-54. doi:10.1136/bjo.2005.072546 pmid:16299149.
- 64 Jaeschke R, Guyatt GH, Sackett DL. Users' guides to the medical literature. III. How to use an article about a diagnostic test. B. What are the results and will they help me in caring for my patients? The Evidence-Based Medicine Working Group. *JAMA* 1994;271:703-7. doi:10.1001/jama.1994.03510330081039 pmid:8309035.
- 65 Davison R, Cannon R. Estimation of central venous pressure by examination of jugular veins. *Am Heart J* 1974;87:279-82. doi:10.1016/0002-8703(74)90064-7 pmid:4812363.
- 66 Connors AF Jr, McCaffree DR, Gray BA. Evaluation of right-heart catheterization in the critically ill patient without acute myocardial infarction. *N Engl J Med* 1983;308:263-7. doi:10.1056/NEJM198302033080508 pmid:6848938.
- 67 Ducas J, Magder S, McGregor M. Validity of the hepatojugular reflux as a clinical test for congestive heart failure. *Am J Cardiol* 1983;52:1299-303. doi:10.1016/0002-9149(83)90592-1 pmid:6650420.
- 68 Cook DJ. Clinical assessment of central venous pressure in the critically ill. *Am J Med Sci* 1990;299:175-8. doi:10.1097/00000441-199003000-00006 pmid:2316561.
- 69 Stein JH, Neumann A, Marcus RH. Comparison of estimates of right atrial pressure by physical examination and echocardiography in patients with congestive heart failure and reasons for discrepancies. *Am J Cardiol* 1997;80:1615-8. doi:10.1016/S0002-9149(97)00776-5 pmid:9416951.
- 70 Vinayak AG, Levitt J, Gehlbach B, Pohlman AS, Hall JB, Kress JP. Usefulness of the external jugular vein examination in detecting abnormal central venous pressure in critically ill patients. *Arch Intern Med* 2006;166:2132-7. doi:10.1001/archinte.166.19.2132 pmid:17060544.
- 71 Sinisalo J, Rapola J, Rossinen J, Kupari M. Simplifying the estimation of jugular venous pressure. *Am J Cardiol* 2007;100:1779-81. doi:10.1016/j.amjcard.2007.07.030 pmid:18082526.
- 72 Brennan JM, Blair JE, Goonewardena S, et al. A comparison by medicine residents of physical examination versus hand-carried ultrasound for estimation of right atrial pressure. *Am J Cardiol* 2007;99:1614-6. doi:10.1016/j.amjcard.2007.01.037 pmid:17531592.
- 73 Deol GR, Collett N, Ashby A, Schmidt GA. Ultrasound accurately reflects the jugular venous examination but underestimates central venous pressure. *Chest* 2011;139:95-100. doi:10.1378/chest.10-1301 pmid:20798190.
- 74 McGee SR. Physical examination of venous pressure: a critical review. *Am Heart J* 1998;136:10-8. doi:10.1016/S0002-8703(98)70175-9 pmid:9665212.
- 75 Seth R, Magner P, Matzinger F, van Walraven C. How far is the sternal angle from the mid-right atrium? *J Gen Intern Med* 2002;17:852-6. doi:10.1046/j.1525-1497.2002.20101.x pmid:12406357.
- 76 Sankoff J, Zidulka A. Non-invasive method for the rapid assessment of central venous pressure: description and validation by a single examiner. *West J Emerg Med* 2008;9:201-5. pmid:19561745.
- 77 Sochowski RA, Dubbin JD, Naqvi SZ. Clinical and hemodynamic assessment of the hepatojugular reflux. *Am J Cardiol* 1990;66:1002-6. doi:10.1016/0002-9149(90)90940-3 pmid:2220606.
- 78 Ewy GA. The abdominojugular test: technique and hemodynamic correlates. *Ann Intern Med* 1988;109:456-60. doi:10.7326/0003-4819-109-6-456 pmid:3415106.
- 79 Butman SM, Ewy GA, Standen JR, Kern KB, Hahn E. Bedside cardiovascular examination in patients with severe chronic heart failure: importance of rest or inducible jugular venous distension. *J Am Coll Cardiol* 1993;22:968-74. doi:10.1016/0735-1097(93)90405-P pmid:8409071.
- 80 Grissom CK, Morris AH, Lanken PN, et al. National Institutes of Health/ National Heart, Lung and Blood Institute Acute Respiratory Distress. Association of physical examination with pulmonary artery catheter parameters in acute lung injury. *Crit Care Med* 2009;37:2720-6. doi:10.1097/CCM.0b013e3181a59532 pmid:19885995.
- 81 Ait-Oufella H, Bige N, Boelle PY, et al. Capillary refill time exploration during septic shock. *Intensive Care Med* 2014;40:958-64. doi:10.1007/s00134-014-3326-4 pmid:24811942.
- 82 Mrgan M, Rytter D, Brabrand M. Capillary refill time is a predictor of short-term mortality for adult patients admitted to a medical department: an observational cohort study. *Emerg Med J* 2014;31:954-8. doi:10.1136/emmed-2013-202925 pmid:24045049.
- 83 Kortbeek BJ, Al Turki AS, Ali J. *Advanced trauma life support for doctors, student course manual*. 8th ed. American College of Surgeons, 2008.
- 84 Dellinger RP, Levy MM, Rhodes A, et al. Surviving Sepsis Campaign Guidelines Committee including the Pediatric Subgroup. Surviving sepsis campaign: international guidelines for management of severe sepsis and septic shock: 2012. *Crit Care Med* 2013;41:580-637. doi:10.1097/CCM.0b013e31827e83af pmid:23353941.
- 85 Fink HA, Lederle FA, Roth CS, Bowles CA, Nelson DB, Haas MA. The accuracy of physical examination to detect abdominal aortic aneurysm. *Arch Intern Med* 2000;160:833-6. doi:10.1001/archinte.160.6.833 pmid:10737283.
- 86 Twomey A, Twomey E, Wilkins RA, Lewis JD. Unrecognised aneurysmal disease in male hypertensive patients. *Int Angiol* 1986;5:269-73. pmid:3295075.
- 87 Allen PI, Gourevitch D, McKinley J, Tudway D, Goldman M. Population screening for aortic aneurysms. *Lancet* 1987;2:736. doi:10.1016/S0140-6736(87)91090-7 pmid:2888954.
- 88 al-Zahrani HA, Rawas M, Maimani A, Gasab M, Aba al Khalil BA. Screening for abdominal aortic aneurysm in the Jeddah area, western Saudi Arabia. *Cardiovasc Surg* 1996;4:87-92. doi:10.1016/0967-2109(96)83791-4 pmid:8634854.
- 89 Andersson AP, Ellitsgaard N, Jorgensen B, et al. Screening for abdominal aortic aneurysm in 295 outpatients with intermittent claudication. *Vasc Surg* 1991;25:516-20. doi:10.1177/153857449102500702.
- 90 MacSweeney ST, O'Meara M, Alexander C, O'Malley MK, Powell JT, Greenhalgh RM. High prevalence of unsuspected abdominal aortic aneurysm in patients with confirmed symptomatic peripheral or cerebral arterial disease. *Br J Surg* 1993;80:582-4. doi:10.1002/bjs.1800800510 pmid:8518892.
- 91 Lederle FA, Simel DL. The rational clinical examination. Does this patient have abdominal aortic aneurysm? *JAMA* 1999;281:77-82. doi:10.1001/jama.281.1.77 pmid:9892455.
- 92 Lederle FA, Walker JM, Reinke DB. Selective screening for abdominal aortic aneurysms with physical examination and ultrasound. *Arch Intern Med* 1988;148:1753-6. doi:10.1001/archinte.1988.00380080049015 pmid:3041938.
- 93 Karkos CD, Mukhopadhyay U, Papakostas I, Ghosh J, Thomson GJ, Hughes R. Abdominal aortic aneurysm: the role of clinical examination and opportunistic detection. *Eur J Vasc Endovasc Surg* 2000;19:299-303. doi:10.1053/ejvs.1999.1002 pmid:10753695.
- 94 Cooke G, Doust J, Sanders S. Is pulse palpation helpful in detecting atrial fibrillation? A systematic review. *J Fam Pract* 2006;55:130-4. pmid:16451780.
- 95 Harris K, Edwards D, Mant J. How can we best detect atrial fibrillation? *R Coll Physicians Edinb* 2012;42(Suppl 18):5-22. pmid:22518390.
- 96 Morgan S, Mant D. Randomised trial of two approaches to screening for atrial fibrillation in UK general practice. *Br J Gen Pract* 2002;52:373-4. 377-80. pmid:12014534.
- 97 Sudlow M, Rodgers H, Kenny RA, Thomson R. Identification of patients with atrial fibrillation in general practice: a study of screening methods. *BMJ* 1998;317:327-8. doi:10.1136/bmj.317.7154.327 pmid:9685281.
- 98 Somerville S, Somerville J, Croft P, Lewis M. Atrial fibrillation: a comparison of methods to identify cases in general practice. *Br J Gen Pract* 2000;50:727-9. pmid:11050790.
- 99 Khan NA, Rahim SA, Anand SS, Simel DL, Panju A. Does the clinical examination predict lower extremity peripheral arterial disease? *JAMA* 2006;295:536-46. doi:10.1001/jama.295.5.536 pmid:16449619.
- 100 Armstrong DW, Tobin C, Matangi MF. The accuracy of the physical examination for the detection of lower extremity peripheral arterial disease. *Can J Cardiol* 2010;26:e346-50. doi:10.1016/S0828-282X(10)70467-0 pmid:21165366.
- 101 Brueseke TJ, Macrino S, Miller JJ. Lack of lower extremity hair not a predictor for peripheral arterial disease. *Arch Dermatol* 2009;145:1456-7. doi:10.1001/archdermatol.2009.310 pmid:20026862.

- 102 von Kodolitsch Y, Schwartz AG, Nienaber CA. Clinical prediction of acute aortic dissection. *Arch Intern Med* 2000;160:2977-82. doi:10.1001/archinte.160.19.2977 pmid:11041906.
- 103 Hagan PG, Nienaber CA, Isselbacher EM, et al. The International Registry of Acute Aortic Dissection (IRAD): new insights into an old disease. *JAMA* 2000;283:897-903. doi:10.1001/jama.283.7.897 pmid:10685714.
- 104 Bossone E, Rampoldi V, Nienaber CA, et al. International Registry of Acute Aortic Dissection (IRAD) Investigators. Usefulness of pulse deficit to predict in-hospital complications and mortality in patients with acute type A aortic dissection. *Am J Cardiol* 2002;89:851-5. doi:10.1016/S0002-9149(02)02198-7 pmid:11909573.
- 105 Grim CE, Luft FC, Weinberger MH, Grim CM. Sensitivity and specificity of screening tests for renal vascular hypertension. *Ann Intern Med* 1979;91:617-22. doi:10.7326/0003-4819-91-4-617 pmid:484965.
- 106 Krijnen P, van Jaarsveld BC, Steyerberg EW, Man in 't Veld AJ, Schalekamp MA, Habbema JD. A clinical prediction rule for renal artery stenosis. *Ann Intern Med* 1998;129:705-11. doi:10.7326/0003-4819-129-9-199811010-00005 pmid:9841602.
- 107 Carmichael DJ, Mathias CJ, Snell ME, Peart S. Detection and investigation of renal artery stenosis. *Lancet* 1986;1:667-70. doi:10.1016/S0140-6736(86)91738-1 pmid:2419718.
- 108 Simon N, Franklin SS, Bleifer KH, Maxwell MH. Clinical characteristics of renovascular hypertension. *JAMA* 1972;220:1209-18. doi:10.1001/jama.1972.03200090031005 pmid:5067309.
- 109 Svetkey LP, Helms MJ, Dunnick NR, Klotman PE. Clinical characteristics useful in screening for renovascular disease. *South Med J* 1990;83:743-7. doi:10.1097/00007611-199007000-00005 pmid:2371594.
- 110 McColgan P, Bentley P, McCarron M, Sharma P. Evaluation of the clinical utility of a carotid bruit. *QJM* 2012;105:1171-7. doi:10.1093/qjmed/hcs140 pmid:22886230.
- 111 Brott TG, Halperin JL, Abbara S, et al. American College of Cardiology Foundation American Stroke Association American Association of Neurological Surgeons American College of Radiology American Society of Neuroradiology Congress of Neurological Surgeons Society of Atherosclerosis Imaging and Prevention Society for Cardiovascular Angiography and Interventions Society of Interventional Radiology Society of NeuroInterventional Surgery Society for Vascular Medicine Society for Vascular Surgery. 2011 ASA/ACCF/AHA/AANN/AANS/ACR/ASNR/CNS/SAIP/SCAI/SIR/SNIS/SVM/SVS guideline on the management of patients with extracranial carotid and vertebral artery disease: executive summary. A report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American Stroke Association, American Association of Neuroscience Nurses, American Association of Neurological Surgeons, American College of Radiology, American Society of Neuroradiology, Congress of Neurological Surgeons, Society of Atherosclerosis Imaging and Prevention, Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, Society of NeuroInterventional Surgery, Society for Vascular Medicine, and Society for Vascular Surgery. *Circulation* 2011;124:489-532. doi:10.1161/CIR.0b013e31820d8d78 pmid:21282505.
- 112 Wang CS, FitzGerald JM, Schulzer M, Mak E, Ayas NT. Does this dyspneic patient in the emergency department have congestive heart failure? *JAMA* 2005;294:1944-56. doi:10.1001/jama.294.15.1944 pmid:16234501.
- 113 Madhok V, Falk G, Rogers A, Struthers AD, Sullivan FM, Fahey T. The accuracy of symptoms, signs and diagnostic tests in the diagnosis of left ventricular dysfunction in primary care: a diagnostic accuracy systematic review. *BMC Fam Pract* 2008;9:56. doi:10.1186/1471-2296-9-56 pmid:18842141.
- 114 Damy T, Kallvikbacka-Bennett A, Zhang J, et al. Does the physical examination still have a role in patients with suspected heart failure? *Eur J Heart Fail* 2011;13:1340-8. doi:10.1093/eurjhf/hfr128 pmid:21990340.
- 115 Kelder JC, Cramer MJ, van Wijngaarden J, et al. The diagnostic value of physical examination and additional testing in primary care patients with suspected heart failure. *Circulation* 2011;124:2865-73. doi:10.1161/CIRCULATIONAHA.111.019216 pmid:22104551.
- 116 Drazner MH, Rame JE, Dries DL. Third heart sound and elevated jugular venous pressure as markers of the subsequent development of heart failure in patients with asymptomatic left ventricular dysfunction. *Am J Med* 2003;114:431-7. doi:10.1016/S0002-9343(03)00058-5 pmid:12727575.
- 117 Drazner MH, Rame JE, Stevenson LW, Dries DL. Prognostic importance of elevated jugular venous pressure and a third heart sound in patients with heart failure. *N Engl J Med* 2001;345:574-81. doi:10.1056/NEJMoA010641 pmid:11529211.
- 118 Damman K, Voors AA, Hillege HL, et al. CIBIS-2 Investigators and Committees. Congestion in chronic systolic heart failure is related to renal dysfunction and increased mortality. *Eur J Heart Fail* 2010;12:974-82. doi:10.1093/eurjhf/hfq118 pmid:20685688.
- 119 Caldentey G, Khairy P, Roy D, et al. Prognostic value of the physical examination in patients with heart failure and atrial fibrillation: insights from the AF-CHF trial (atrial fibrillation and chronic heart failure). *JACC Heart Fail* 2014;2:15-23. doi:10.1016/j.jchf.2013.10.004 pmid:24622114.
- 120 Minami Y, Kajimoto K, Sato N, et al. Third heart sound in hospitalised patients with acute heart failure: insights from the ATTEND study. *Int J Clin Pract* 2015;69:820-8. doi:10.1111/ijcp.12603 pmid:25521285.
- 121 Lala A, McNulty SE, Mentz RJ, et al. Relief and Recurrence of Congestion During and After Hospitalization for Acute Heart Failure: Insights From Diuretic Optimization Strategy Evaluation in Acute Decompensated Heart Failure (DOSE-AHF) and Cardiorenal Rescue Study in Acute Decompensated Heart Failure (CARESS-HF). *Circ Heart Fail* 2015;8:741-8. doi:10.1161/CIRCHEARTFAILURE.114.001957 pmid:26041600.
- 122 Lucas C, Johnson W, Hamilton MA, et al. Freedom from congestion predicts good survival despite previous class IV symptoms of heart failure. *Am Heart J* 2000;140:840-7. doi:10.1067/mhj.2000.110933 pmid:11099986.
- 123 Ambrosy AP, Pang PS, Khan S, et al. EVEREST Trial Investigators. Clinical course and predictive value of congestion during hospitalization in patients admitted for worsening signs and symptoms of heart failure with reduced ejection fraction: findings from the EVEREST trial. *Eur Heart J* 2013;34:835-43. doi:10.1093/eurheartj/ehs444 pmid:23293303.
- 124 McMurray JJ, Adamopoulos S, Anker SD, et al. ESC Committee for Practice Guidelines. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: The Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur Heart J* 2012;33:1787-847. doi:10.1093/eurheartj/ehs104 pmid:22611136.
- 125 Yancy CW, Jessup M, Bozkurt B, et al. WRITING COMMITTEE MEMBERS American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation* 2013;128:e240-327. doi:10.1161/CIR.0b013e31829e8807 pmid:23741058.
- 126 Fahey T, Jeyaseelan S, McCowan C, et al. Diagnosis of left ventricular systolic dysfunction (LVSD): development and validation of a clinical prediction rule in primary care. *Fam Pract* 2007;24:628-35. doi:10.1093/fampra/cmm055 pmid:17827466.
- 127 Etchells E, Bell C, Robb K. Does this patient have an abnormal systolic murmur? *JAMA* 1997;277:564-71. doi:10.1001/jama.1997.03540310062036 pmid:9032164.
- 128 Douglas PS, Garcia MJ, Haines DE, et al. American College of Cardiology Foundation Appropriate Use Criteria Task Force American Society of Echocardiography American Heart Association American Society of Nuclear Cardiology Heart Failure Society of America Heart Rhythm Society Society for Cardiovascular Angiography and Interventions Society of Critical Care Medicine Society of Cardiovascular Computed Tomography Society for Cardiovascular Magnetic Resonance American College of Chest Physicians. ACCF/AHA/ASA/ASNC/HFSA/HRS/SCAI/SCCM/SCCT/SCMR 2011 Appropriate Use Criteria for Echocardiography. A Report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, American Society of Echocardiography, American Heart Association, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Critical Care Medicine, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance American College of Chest Physicians. *J Am Soc Echocardiogr* 2011;24:229-67. pmid:21338862.
- 129 British Society of Echocardiography. Clinical indications for echocardiography. http://www.bsecho.org/media/64844/indications_for_echocardiography.pdf.
- 130 Attenhofer Jost CH, Turina J, Mayer K, et al. Echocardiography in the evaluation of systolic murmurs of unknown cause. *Am J Med* 2000;108:614-20. doi:10.1016/S0002-9343(00)00361-2 pmid:10856408.
- 131 Mishra M, Chambers JB, Jackson G. Murmurs in pregnancy: an audit of echocardiography. *BMJ* 1992;304:1413-4. doi:10.1136/bmj.304.6839.1413 pmid:1628016.
- 132 Ahuja IM. Functional systolic murmurs. *Indian Heart J* 1982;34:241-4. pmid:7141453.
- 133 Lockhart PB, Crist D, Stone PH. The reliability of the medical history in the identification of patients at risk for infective endocarditis. *J Am Dent Assoc* 1989;119:417-8. 421-2. doi:10.14219/jada.archive.1989.0060 pmid:2527900.
- 134 Roldan CA, Shively BK, Crawford MH. Value of the cardiovascular physical examination for detecting valvular heart disease in asymptomatic subjects. *Am J Cardiol* 1996;77:1327-31. doi:10.1016/S0002-9149(96)00200-7 pmid:8677874.
- 135 Reichlin S, Dieterle T, Camli C, Leimenstoll B, Schoenenberger RA, Martina B. Initial clinical evaluation of cardiac systolic murmurs in the ED by noncardiologists. *Am J Emerg Med* 2004;22:71-5. doi:10.1016/S0735-6757(03)00093-7 pmid:15011216.
- 136 Aronow WS, Kronzon I. Correlation of prevalence and severity of valvular aortic stenosis determined by continuous-wave Doppler echocardiography with physical signs of aortic stenosis in patients aged 62 to 100 years with aortic systolic ejection murmurs. *Am J Cardiol* 1987;60:399-401. doi:10.1016/0002-9149(87)90262-1 pmid:3497570.

- 137 Etchells E, Glens V, Shadowitz S, Bell C, Siu S. A bedside clinical prediction rule for detecting moderate or severe aortic stenosis. *J Gen Intern Med* 1998;13:699-704. doi:10.1046/j.1525-1497.1998.00207.x pmid:9798818.
- 138 Loxdale SJ, Sneyd JR, Donovan A, Werrett G, Viira DJ. The role of routine pre-operative bedside echocardiography in detecting aortic stenosis in patients with a hip fracture. *Anaesthesia* 2012;67:51-4. doi:10.1111/j.1365-2044.2011.06942.x pmid:22023667.
- 139 Lehmann KG, Francis CK, Dodge HT. TIMI Study Group. Mitral regurgitation in early myocardial infarction. Incidence, clinical detection, and prognostic implications. *Ann Intern Med* 1992;117:10-7. doi:10.7326/0003-4819-117-1-10 pmid:1596042.
- 140 Panidis IP, McAllister M, Ross J, Mintz GS. Prevalence and severity of mitral regurgitation in the mitral valve prolapse syndrome: a Doppler echocardiographic study of 80 patients. *J Am Coll Cardiol* 1986;7:975-81. doi:10.1016/S0735-1097(86)80214-5 pmid:3958380.
- 141 Rahko PS. Prevalence of regurgitant murmurs in patients with valvular regurgitation detected by Doppler echocardiography. *Ann Intern Med* 1989;111:466-72. doi:10.7326/0003-4819-111-6-466 pmid:2774371.
- 142 Maisel AS, Atwood JE, Goldberger AL. Hepatojugular reflux: useful in the bedside diagnosis of tricuspid regurgitation. *Ann Intern Med* 1984;101:781-2. doi:10.7326/0003-4819-101-6-781 pmid:6497192.
- 143 Choudhry NK, Etchells EE. The rational clinical examination. Does this patient have aortic regurgitation? *JAMA* 1999;281:2231-8. doi:10.1001/jama.281.23.2231 pmid:10376577.
- 144 Grayburn PA, Smith MD, Handshoe R, Friedman BJ, DeMaria AN. Detection of aortic insufficiency by standard echocardiography, pulsed Doppler echocardiography, and auscultation. A comparison of accuracies. *Ann Intern Med* 1986;104:599-605. doi:10.7326/0003-4819-104-5-599 pmid:3963660.
- 145 Aronow WS, Kronzon I. Correlation of prevalence and severity of aortic regurgitation detected by pulsed Doppler echocardiography with the murmur of aortic regurgitation in elderly patients in a long-term health care facility. *Am J Cardiol* 1989;63:128-9. doi:10.1016/0002-9149(89)91098-9 pmid:2491772.
- 146 Alexander WD, Polak A. Early diastolic murmurs in end-stage renal failure. *Br Heart J* 1977;39:900-2. doi:10.1136/hrt.39.8.900 pmid:901686.
- 147 Tribouilloy CM, Enriquez-Sarano M, Mohty D, et al. Pathophysiologic determinants of third heart sounds: a prospective clinical and Doppler echocardiographic study. *Am J Med* 2001;111:96-102. doi:10.1016/S0002-9343(01)00769-0 pmid:11498061.
- 148 Mookherjee S, Pheatt L, Ranji SR, Chou CL. Physical examination education in graduate medical education--a systematic review of the literature. *J Gen Intern Med* 2013;28:1090-9. doi:10.1007/s11606-013-2380-x pmid:23568186.
- 149 Rothberg MB. A piece of my mind. The \$50,000 physical. *JAMA* 2014;311:2175-6. doi:10.1001/jama.2014.3415 pmid:24893085.
- 150 Bergl P, Farnan JM, Chan E. Moving toward cost-effectiveness in physical examination. *Am J Med* 2015;128:109-10. doi:10.1016/j.amjmed.2014.10.003 pmid:25305232.
- 151 Elder A, Chi J, Ozdalga E, Kugler J, Verghese A. A piece of my mind. The road back to the bedside. *JAMA* 2013;310:799-800. doi:10.1001/jama.2013.227195 pmid:23982364.
- 152 National Academies of Sciences, Engineering, and Medicine. *Improving diagnosis in health care*. National Academies Press, 2015.
- 153 Herrle SR, Corbett EC Jr, Fagan MJ, Moore CG, Elnicki DM. Bayes' theorem and the physical examination: probability assessment and diagnostic decision making. *Acad Med* 2011;86:618-27. doi:10.1097/ACM.0b013e318212eb00 pmid:21436660.
- 154 Barrett PM, Topol EJ. To truly look inside. *Lancet* 2016;387:1268-9. doi:10.1016/S0140-6736(16)30027-7 pmid:27035018.