

High rates of clinically relevant incidental findings by total-body CT scanning in trauma patients; results of the REACT-2 trial

K. Treskes¹ · S. A. Bos¹ · L. F. M. Beenen² · J. C. Sierink¹ · M. J. R. Edwards³ ·
B. J. A. Beuker⁴ · G. S. R. Muradin⁵ · J. Hohmann⁶ · J. S. K. Luitse¹ · M. W. Hollmann⁷ ·
M. G. W. Dijkgraaf⁸ · J. C. Goslings¹ · on behalf of the REACT-2 study group

Received: 5 April 2016 / Revised: 21 July 2016 / Accepted: 5 September 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract

Objectives To determine whether there is a difference in frequency and clinical relevance of incidental findings detected by total-body computed tomography scanning (TBCT) compared to those by the standard work-up (STWU) with selective computed tomography (CT) scanning.

Presentations Abstract presented at the annual meeting of the Dutch society for trauma surgery and Dutch orthopaedic society (November 5, 2015; Amsterdam, The Netherlands)

✉ K. Treskes
k.treskes@amc.nl

- ¹ Trauma Unit, Department of Surgery, Academic Medical Center, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands
- ² Department of Radiology, Academic Medical Center, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands
- ³ Department of Trauma and emergency surgery, Radboud University Medical Center, Geert Grooteplein-Zuid 10, 6525 GA Nijmegen, The Netherlands
- ⁴ Trauma Unit, Department of Surgery, University Medical Center Groningen, Hanzeplein 1, 9700 RB Groningen, The Netherlands
- ⁵ Department of Radiology, Erasmus MC, University Medical Center Rotterdam, 's-Gravendijkwal 230, 3015 CE Rotterdam, The Netherlands
- ⁶ Department of Radiology and Nuclear Medicine, University of Basel Hospital, Petersgraben 4, 4031 Basel, Switzerland
- ⁷ Department of Anaesthesiology, Academic Medical Center, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands
- ⁸ Clinical Research Unit, Academic Medical Center, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands

Methods Trauma patients from five trauma centres were randomized between April 2011 and January 2014 to TBCT imaging or STWU consisting of conventional imaging with selective CT scanning. Incidental findings were divided into three categories: 1) major finding, may cause mortality; 2) moderate finding, may cause morbidity; and 3) minor finding, hardly relevant. Generalized estimating equations were applied to assess differences in incidental findings.

Results In total, 1083 patients were enrolled, of which 541 patients (49.9 %) were randomized for TBCT and 542 patients (50.1 %) for STWU. Major findings were detected in 23 patients (4.3 %) in the TBCT group compared to 9 patients (1.7 %) in the STWU group (adjusted rate ratio 2.851; 95%CI 1.337–6.077; $p < 0.007$). Findings of moderate relevance were detected in 120 patients (22.2 %) in the TBCT group compared to 86 patients (15.9 %) in the STWU group (adjusted rate ratio 1.421; 95%CI 1.088–1.854; $p < 0.010$).

Conclusions Compared to selective CT scanning, more patients with clinically relevant incidental findings can be expected by TBCT scanning.

Key points

- Total-body CT scanning in trauma results in 1.5 times more incidental findings.
- Evaluation by TBCT in trauma results in more patients with incidental findings.
- In every category of clinical relevance, TBCT detects more incidental findings.

Keywords Total-body CT · Incidental findings · Multiple trauma · Wounds and injuries · Multidetector computed tomography

Abbreviations

CT	Computed tomography
TBCT	Total-body computed tomography
STWU	Standard work-up
FAST	Focused assessment with sonography for trauma
GEE	Generalized estimating equation

Introduction

Total-body computed tomography scanning (TBCT) is often used during the primary assessment of patients after severe trauma. Instead of selective computed tomography (CT) scanning of specific body regions, trauma teams routinely perform CT scans of the head, neck, thorax, abdomen and pelvis for trauma patients who could benefit from TBCT scanning. A potential disadvantage compared to the selective approach is the increased radiation exposure of TBCT scanning [1, 2]. Since TBCT has not been shown to reduce mortality in the general trauma population, indication setting is important and a subject of debate [3]. Another consideration when performing TBCT is the increase in non-trauma-related radiologic findings. These concomitant incidental findings should be prioritized with respect to potential life-threatening injuries, and may require additional follow-up and treatment. Incidental findings could provide the advantage of earlier diagnosis of malignancy or vascular disease. On the contrary, when clinical significance is absent, incidental findings could also result in unnecessary investigations and concerns for the patient and extra health care costs.

Previous studies have reported the detection of incidental findings in 32 % to 43 % of trauma patients screened in trauma centres with selective CT scanning [4, 5], and studies on TBCT scanning have reported incidental findings in 45 % to 53 % of trauma patients [6–8]. However, direct comparisons of the frequency and relevance of incidental findings between TBCT and standard work-up with selective CT scanning in trauma patients are lacking. A difference in the frequency of relevant incidental findings can therefore only be assumed.

The aim of this study was to determine whether incidental findings detected by TBCT scanning differed in frequency and relevance from those detected by conventional imaging supplemented with selective CT in patients with severe trauma.

Patients and methods

Study design and patient selection

This study was a secondary analysis of patients selected for the REACT-2 multicentre randomized controlled trial, of which the study protocol and main results were published previously [3, 9]. In short, in the REACT-2 trial, adult trauma

patients with compromised vital parameters, clinical suspicion of specific severe injuries or high-risk trauma mechanisms were randomized to undergo either an immediate TBCT scan or standard radiological work-up (STWU). The inclusion and exclusion criteria are listed in the table 4 in the appendix. Patients included those seen between April 2011 and January 2014 in four level 1 trauma centres in the Netherlands and one in Switzerland. Informed consent was temporarily waived during the initial presentation in the trauma room. At the earliest opportunity after the trauma, patients or their legal representatives were given work-up information and informed consent was requested. The study was approved by the medical ethics committees at all participating centres (AMC MEC 10/145).

The randomization process was performed by the trauma team immediately after the primary assessment of the patient. TBCT scanning was performed without conventional imaging or sonography in advance and consisted of a non-enhanced CT scan of the head and neck, with arms alongside the trunk, followed by a contrast-enhanced CT scan of the chest, abdomen and pelvis. The preferred technique for the second scan was split-bolus intravenous contrast imaging after raising of the arms, if possible [10]. STWU consisted of x-rays of the chest and pelvis, a focused assessment with sonography for trauma (FAST), and CT scans from specific body regions if indicated. Indications for selective CT scanning were predefined according to local protocols. These indications are listed in the table 4 in the appendix. CT scanners were located in the trauma room or in an adjacent room, and all were 64-slice multidetector row CT scanners.

Data collection

Radiological images were interpreted by the radiology resident and subsequently by a senior radiologist experienced in trauma imaging. Although focusing on traumatic injuries, this ‘double-reading system’ minimizes the number of missed findings [11]. All findings were described in the radiological reports, which are accessible through the computerized hospital databases of participating centres. Any available previous radiologic imaging of the same patient was also reviewed to confirm the findings to be new. In addition, trauma room reports, interventional and pathology reports, and discharge letters were reviewed. For follow-up data, all available and relevant in-hospital files were searched within a minimum of 6 months and maximum of 2 years after admission to the trauma resuscitation room.

Definitions and categorization of incidental findings

The clinical relevance of an incidental finding was divided into three subcategories: 1) major finding, may cause mortality; 2) moderate finding, may cause morbidity; and 3) minor finding, hardly relevant and no follow-up needed. The

findings and corresponding relevance were scored based on the latest information on the finding. Using earlier reports on this subject, a list of incidental findings that could be expected was formulated before data acquisition [4–8, 12, 13]. Some incidental findings were added to more than one relevance category, as the clinical importance of the same type of finding varied widely. For these specific findings, size-, age- or complexity-specific cut-off values have been added to their description (e.g. simple vs. complicated renal cyst). Findings that had been described previously, as well as traumatic lesions, were excluded. Degenerative joint diseases, common atherosclerotic vessel disease, enostosis, sinusitis, age-related cerebral atrophy, and signs of earlier operations or old cerebral hematoma/infarction were also excluded as incidental findings, in accordance with previous literature [4, 8, 14].

For pulmonary nodules and renal cysts, classification was performed in accordance with the Fleischner society pulmonary nodule recommendations and Bosniak renal cyst classification [15, 16]. With respect to the Fleischner recommendations, findings in patients requiring follow-up within 6 months were defined as major findings, follow-up between 6 and 12 months as moderate findings, and minor if no follow-up was needed. Bosniak class 1 corresponds to minor findings, class 2 to moderate findings, and classes 2 F and over to major findings. In the case of any abnormal lymph nodes, asymptomatic findings were classified as minor unless the nodes were >10 mm in size, in which case they were labelled as moderate. Lymphadenopathy of major relevance indicates suspected lethal findings such as a suspected (non-)Hodgkin lymphoma. If the nodular size was not reported, classification was performed by the reviewers of this study according to follow-up advice or further descriptions from the reporting radiologist.

Multitrauma patients were defined as patients with an Injury Severity Score ≥ 16 . Traumatic Brain Injury (TBI) patients were defined as patients with a Glasgow Coma Scale (GCS) score <9 at presentation and Abbreviated Injury Scale (AIS)-Head code ≥ 3 .

Statistical analysis

Continuous data with a normal distribution are presented as means with standard deviation, and the non-normally distributed data are presented as medians with interquartile ranges (IQR). Independent samples *t* tests and Mann–Whitney *U* tests were used to compare the parametric and non-parametric continuous data, respectively. Generalized estimating equations (GEE) were applied to assess differences in incidental findings between TBCT and STWU. The Kolmogorov–Smirnov test was used to confirm Poisson-distributed numbers of major and moderate incidental findings. The total number of incidental findings, as well as the number of minor incidental findings, seemed to be negative

binomially distributed, which was confirmed by a non-significant chi-square test of goodness of fit of observed and theoretical data, the latter generated with the ‘rnegbin’ function in the R computing environment. The results are reported as rate ratios of incidental findings with TBCT versus STWU, corrected for age, sex and centre. The chi-square test was used to compare categorical variables when categories consisted of at least ten cases; otherwise, Fisher’s exact test was used. A *p* value below 0.05 was considered to reflect statistical significance. Statistical analysis was performed with SPSS version 23 software (IBM Corp., Armonk, NY, USA).

Results

Baseline characteristics of the study population are shown in Table 1. In total, 1083 patients were enrolled, of which 541 patients were randomized for TBCT ($n = 49.9\%$). The median age was 42 years (IQR 27–59) in the TBCT group and 45 years (IQR 26–59) in the STWU group ($p = 0.746$). The baseline characteristics were comparable between groups except for the number of multitrauma patients: TBCT $n = 362$ (66.9 %) vs. STWU $n = 331$ (61.1 %), $p = 0.045$. Median ISS did not differ between groups: TBCT 20 (IQR 10–29) vs. STWU 19 (9–29), $p = 0.405$.

In total, 441 incidental findings were found in 233 of the patients (42.9 %) randomized for TBCT, compared to 290 findings in 167 of the patients (32.5 %) randomized for the STWU (adjusted rate ratio 1.531; 95 % confidence interval [95%CI] 1.274–1.840; $p < 0.001$), as shown in Table 2. Major findings were detected in 23 patients (4.3 %) in the TBCT group, compared to 9 patients (1.7 %) in the STWU group (adjusted rate ratio 2.851; 95%CI 1.337–6.077; $p < 0.007$). Moderate findings were detected in 120 patients (22.2 %) and 86 patients (15.9 %) in the TBCT and STWU groups, respectively (adjusted rate ratio 1.421; 95%CI 1.088–1.854; $p < 0.010$).

Table 3 shows comparisons of the distribution of the incidental findings over clinical categories between the two groups. Distribution over categories of relevance, body regions, organ systems or neoplasms was similar between the imaging groups. Tables 5 and 6 in the appendix show the follow-up and medical documentation of incidental findings per category of relevance. These characteristics were comparable between the imaging groups; however, follow-up rates were low, and documentation of incidental findings was poor in both groups. In the discharge letters, 39.3 % of the major findings and 13.8 % of the moderate findings were mentioned.

The complete list of all findings arranged by body region and relevance is presented in Table 7 in the appendix. Simple renal and hepatic cysts were the most common of all incidental findings for patients in both imaging groups. Suspicious

Table 1 Baseline demographic and clinical characteristics of patients*

Characteristic	Total-body CT (<i>n</i> = 541)	Standard work-up (<i>n</i> = 542)
Age (years)	42 (27–59)	45 (26–59)
Male sex, <i>n</i> (%)	413 (76.3)	411 (75.8)
Blunt trauma, <i>n</i> (%)	530 (98.0)	534 (98.5)
Trauma mechanism blunt trauma, <i>n</i> (%)		
Fall from height	170 (32.1)	178 (33.3)
MVC – patient as occupant	201 (37.9)	190 (35.6)
MVC – patient as cyclist	65 (12.3)	60 (11.2)
MVC – patient as pedestrian	29 (5.5)	45 (8.4)
Other	65 (12.3)	61 (11.4)
Comorbidity, <i>n</i> (%)		
ASA I or II	495 (95.7)	501 (96.2)
ASA III, IV or V	22 (4.3)	20 (3.8)
CT performed at ED, <i>n</i> (%)*		
Head	539 (99.6)	483 (89.1)
Neck	535 (98.9)	480 (88.6)
Chest	529 (97.8)	315 (58.1)
Abdomen/pelvis	528 (97.6)	278 (51.3)
Abbreviated Injury Scale ≥ 3 , <i>n</i> (%)		
Head	247 (45.7)	218 (40.2)
Chest	229 (42.3)	206 (38.0)
Abdomen/pelvic content	49 (9.1)	67 (12.4)
Pelvis and extremities	150 (27.7)	154 (28.4)
Injury Severity Score (points)	20 (10–29)	19 (9–29)
Multitrauma patients, <i>n</i> (%) [†]	362 (66.9)	331 (61.1)
TBI patients, <i>n</i> (%) [†]	178 (32.9)	151 (27.9)

* Results in this table were published earlier [3]. $p > 0.05$ for all between-group comparisons except for CT performed ($p < 0.001$ for all body regions) and multitrauma patients ($p = 0.045$). All data are number (%) or median (interquartile range)

[†] Multitrauma patients are defined as ISS ≥ 16 . Traumatic Brain Injury (TBI) patients are defined as GCS < 9 (Glasgow Coma Scale) at presentation and Abbreviated Injury Scale (AIS)-Head ≥ 3

MVC Motor vehicle collision, ASA American Society of Anaesthesiologists, ED emergency department

pulmonary nodules were the most described potentially lethal finding ($n = 6$). Of all findings of moderate relevance, gallstones and hepatic steatosis were most frequently described. One in every 24 incidental findings was a pathologically confirmed neoplasm (4.1 %).

Discussion

This study shows that TBCT imaging is more likely to detect an incidental finding than the standard work-up with selective CT scanning. In every category of clinical relevance, the

Table 2 Trauma patients with incidental findings

Characteristic	Total-body CT (<i>n</i> = 541)	Standard work-up (<i>n</i> = 542)	Rate ratio (95%CI)	<i>p</i>	Adjusted rate ratio* (95%CI)	<i>p</i>
Patients with incidental findings, <i>n</i> (%)	232 (42.9)	176 (32.5)	1.524 (1.251–1.856)	$< 0.001^{\dagger}$	1.531 (1.274–1.840)	$< 0.001^{\dagger}$
Major relevance	23 (4.3)	9 (1.7)	2.672 (1.243–5.744)	0.012 [‡]	2.851 (1.337–6.077)	0.007 [‡]
Moderate relevance	120 (22.2)	86 (15.9)	1.394 (1.051–1.849)	0.021 [‡]	1.421 (1.088–1.854)	0.010 [‡]
Minor relevance	172 (31.8)	129 (23.8)	1.551 (1.240–1.940)	$< 0.001^{\dagger}$	1.536 (1.238–1.905)	$< 0.001^{\dagger}$

*Adjusted for age, sex and centre, [†] Wald test with assumption of negative binomial distribution, [‡] Wald test with assumption of Poisson distribution
95 %CI = 95 % confidence interval

Table 3 Characteristics of incidental findings

Characteristic	Total-body CT	Standard work-up	<i>p</i> *
Incidental findings, <i>n</i>	441	290	
Body region, <i>n</i> (%)			
Head	56 (12.7)	39 (13.4)	0.786
Neck	26 (5.9)	22 (7.6)	
Thorax	65 (14.7)	47 (16.2)	
Abdomen/pelvis	292 (66.2)	180 (62.1)	
Extremities	2 (0.5)	2 (0.7)	
Organ system, <i>n</i> (%)			
Reno-adrenal	112 (25.4)	75 (25.9)	0.696
Hepatobiliary	92 (20.9)	64 (22.1)	
Respiratory	55 (12.5)	27 (9.3)	
Reticuloendothelial	49 (11.1)	33 (11.4)	
Neurological	33 (7.5)	25 (8.6)	
Endocrine	27 (6.1)	21 (7.2)	
Gastrointestinal	27 (6.1)	11 (3.8)	
Urethro-genital	18 (4.1)	15 (5.2)	
Cardiovascular	16 (3.6)	15 (5.2)	
Musculoskeletal	11 (2.5)	4 (1.4)	
Cutaneous	1 (0.2)	0 (0)	
Neoplasm			
Confirmed	20 (4.5)	10 (3.4)	0.456
Suspected	29 (6.6)	14 (4.8)	

* Chi-square test for distribution across categories

TBCT scan detected significantly more findings. The incidental findings did not differ in distribution over body regions or tissue types, although the abdominal region showed the largest difference. We could not demonstrate a significant difference in follow-up, which could be explained by low follow-up rates in general and poor documentation of incidental findings and their management in trauma patients. Trauma teams using TBCT scanning should be aware of an increase in relevant incidental findings and should pay special attention to the reporting and management of these additional findings.

In the present study, incidental findings occurred in 43 % of patients undergoing TBCT scanning, of which 42 % could cause serious morbidity. Similar results were reported in previous studies on incidental findings in TBCT scans for initial trauma evaluation. These studies, however, did not directly compare these to incidental findings with selective CT scanning. The study by Hofstetter et al. found incidental findings in 50 % of their patients, with 29 % of these possibly requiring follow-up [7]. In a study by Barrett et al., findings were detected in 53 % of all patients by TBCT, 59 % of which required urgent follow-up [6]. Sierink et al. recently reported incidental findings in 45 % of all patients, with possible follow-up required in 31 % [8]. Thus, with the present study included, the percentage of trauma patients with incidental findings detected by TBCT ranges from 43 % to 53 %. Of these findings,

29 % to 59 % have clinical relevance; however, the definition of clinical relevance will be interpreted differently.

Considering future diagnostic work-up, the separate and detailed inclusion of moderate and major findings in the trauma room report's conclusion may help communicate these findings to the general practitioner and other treating physicians. Poor documentation could result in lack of further diagnostic work-up or treatment in other facilities, a structural problem that has been described in previous reports on CT in trauma and emergency imaging [4, 5, 13, 17–20]. Complete and clear documentation thus might save costs in the long term by eliminating repeated work-up for incidental findings. Future research should aim to establish effective methods for proper reporting and management of incidental findings in trauma patients.

Limitations and strengths

One limitation of the present study is that the categorization of incidental findings into three relevance groups was subject to personal interpretation, as there is no consensus guideline. Discrepancies between previous studies show that specific findings are not always classified in the same category of clinical relevance. To minimize the effect of interpretation, the categorization of expected incidental findings was performed before data acquisition, in accordance with previous literature and under the supervision of an experienced radiologist. The classification system used in this study closely resembled those of previous studies [4, 6–8, 14].

Second, the documentation of incidental findings in the radiology reports could be incomplete. The number of incidental findings may have been influenced by the acute setting of trauma care, and therefore findings of minor or moderate interest may not have been reported at all, since they seemed irrelevant during primary trauma care. However, the risk of underestimation was reduced by the double-reading system. On the other hand, the rate of unknown findings might be overestimated, because previous imaging of the patient might not have been available during formulation of the radiology report.

Third, the follow-up is likely underestimated due to reporting issues as well. Follow-up was between 6 months and 2 years after the first trauma presentation, and only within the in-hospital documentation of the trauma centres where the patient was initially seen. Subsequently, some patients—for example, those with pulmonary nodules—would receive their first follow-up after 1 year or in a different hospital. Furthermore, it is possible that the finding was discussed and that a watchful waiting approach was preferred but not reported in the patient's file.

This study, which investigated the frequency and clinical relevance of incidental findings in trauma, is the first to directly compare TBCT scanning with conventional imaging supplemented with selective CT. Additional strengths include the international multicentre setting, large comparable patient groups and

its randomization setting. Lastly, the list of expected incidental findings aided in adequate prospective categorization.

Conclusion

When using TBCT instead of selective CT scanning in primary trauma care, a greater number of clinically relevant incidental findings can be expected. Data did not show a significantly higher workload through follow-up; however, documentation of follow-up was suboptimal. When evaluating trauma patients with TBCT scanning, extra alertness towards detection, documentation and follow-up of relevant incidental findings is warranted.

Acknowledgments The scientific guarantor of this publication is Prof. J.C. Goslings. The authors of this manuscript declare no relationships with any companies whose products or services may be related to the subject matter of the article. The REACT-2 has received funding by ZonMw, the Netherlands Organisation for Health Research and Development (grant number: 171102023); Dr. M.G.W. Dijkgraaf has significant statistical expertise. Institutional Review Board approval was obtained. Written informed consent was obtained from all subjects (patients) in this study. All study subjects will be reported in another article reporting the main outcome measures of the REACT-2 trial. [Sierink JC, Treskes K, Edwards MJ et al., Immediate total-body CT scanning versus conventional imaging and selective CT scanning in severe trauma patients: A randomised controlled trial (REACT-2 trial), accepted for publication]. Methodology: prospective, randomised controlled trial, multicentre study.

Collaborators

TP Saltzherr¹, MD, PhD; T Schepers¹, MD, PhD, VM de Jong¹, MD; R van Vugt³, MD, PhD; M Brink⁹, MD, PhD; J Peters³, MD; M El Moumni⁴, MD; JS Harbers⁴, MD; KW Wendt⁴, MD; EMM van Lieshout¹⁰, MSc PhD; MJ Elzinga¹⁰, MD; EH Jansen¹¹, MD; C Zähringer⁶, MD; N Bless¹², MD; R Bingisser¹³, MD

¹ Trauma Unit, Department of Surgery, Academic Medical Center, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands

² Department of Radiology, Academic Medical Center, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands

³ Department of Trauma and emergency surgery, Radboud University Medical Center, Geert Grooteplein-Zuid 10, 6525 GA Nijmegen, The Netherlands

⁴ Trauma Unit, Department of Surgery, University Medical Center Groningen, Hanzeplein 1, 9700 RB Groningen, The Netherlands

⁵ Department of Radiology, Erasmus MC, University Medical Center Rotterdam, 's-Gravendijkwal 230, 3015 CE Rotterdam, The Netherlands

⁶ Department of Radiology and Nuclear Medicine, University of Basel Hospital, Petersgraben 4 4031 Basel, Switzerland

⁷ Department of Anaesthesiology, Academic Medical Center, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands

⁸ Clinical Research Unit, Academic Medical Center, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands

⁹ Department of Radiology, Radboud University Medical Center, Geert Grooteplein-Zuid 10, 6525 GA Nijmegen, The Netherlands

¹⁰ Trauma Research Unit Department of Surgery, Erasmus MC, University Medical Center Rotterdam, 's-Gravendijkwal 230, 3015 CE Rotterdam, The Netherlands

¹¹ Accident and Emergency Department, Erasmus MC, University Medical Center Rotterdam, 's-Gravendijkwal 230, 3015 CE Rotterdam, The Netherlands

¹² Department of Traumatology, University of Basel Hospital, Petersgraben 4, 4031 Basel, Switzerland

¹³ Emergency Unit, University of Basel Hospital, Petersgraben 4, 4031 Basel, Switzerland

Appendix 1 Indications for immediate total-body CT in trauma patients used in REACT-2 trial

Trauma patients with one of the following parameters at hospital arrival:

- respiratory rate ≥ 30 /min or ≤ 10 /min
- pulse ≥ 120 /min
- systolic blood pressure ≤ 100 mmHg
- estimated exterior blood loss ≥ 500 ml
- Glasgow Coma Score ≤ 13
- abnormal pupillary reaction

OR

Patients with a clinical suspicion of one of the following diagnoses:

- fractures from at least two long bones
- flail chest, open chest or multiple rib fractures
- severe abdominal injury
- pelvic fracture
- unstable vertebral fractures/spinal cord compression

OR

Patients with one of the following injury mechanisms:

- fall from a height (>3 meters/ >10 feet)
- ejection from a vehicle
- death of occupant in same vehicle
- severely injured patient in same vehicle
- wedged or trapped chest/abdomen

Contraindications

Trauma patients with any of the following characteristics will be excluded:

- known age < 18 years
- known pregnancy
- referred from another hospital
- clearly low-energy trauma with blunt injury mechanism
- any patient with a stab wound in one body region
- any patient who is judged to be too unstable to undergo a CT scan and requires (cardiopulmonary) resuscitation or immediate operation because death is imminent

Appendix 2

Table 4 Indications for selective CT scanning after conventional imaging

<p>CT-brain</p> <p>A patient with trauma of the head and with at least: → 1 major criterion:</p> <ul style="list-style-type: none"> - EMV \leq13 - loss of consciousness >30 min - haemodynamically unstable - age \geq60 years - high-risk trauma - vomiting - posttraumatic seizure - coagulopathy risk factors (primary or by medication) - focal neurological deficit - >1 point decline in EMV after 1 h - posttraumatic amnesia >4 h - clinical suspicion for skull base or facial fractures <p>→ and/or at least 2 minor criteria:</p> <ul style="list-style-type: none"> - age between 40-60 years - posttraumatic loss of consciousness - posttraumatic amnesia 2-4 h - externally facial injuries without signs of fractures - 1-point decline in EMV after 1 h <p>CT of the cervical spine</p> <ol style="list-style-type: none"> 1. Always when CT-brain is performed 2. In all patients unless they meet all the Nexus criteria: <ul style="list-style-type: none"> - no posterior midline cervical spine tenderness - no focal neurological deficit - a normal level of alertness - no evidence of intoxication - no painful distracting injuries <p>X-cervical spine</p> <p>Never indicated. If Nexus deviant: cervical-CT.</p> <p>Chest CT (with iv contrast)</p> <ol style="list-style-type: none"> 1. Chest gunshot wound with suspicion of transmediastinal route 2. Acute aortic injury 3. Abnormal mediastinum seen at chest radiography. <ul style="list-style-type: none"> - mediastinal widening - pleural cap ('apical cap') - aorta arc unclear enclosed - left main bronchus removed downwards - deviated trachea or oesophagus - filled aortopulmonary window - widened paraspinal line - widened paratracheal line right 3. Relative indications: <ul style="list-style-type: none"> - type and severity of trauma - fractures of costa 1 or 2 - thoracic spine fracture - posterior sternoclavicular luxation - hesitation about the existence of pneumothorax/ pneumomediastinum or pneumopericardium - fractures of the clavicle and shoulder 	<p>Abdominal CT (with iv contrast)</p> <ol style="list-style-type: none"> 1. Penetrating injuries in abdomen, chest and/or flank 2. Deficits found with FAST <ul style="list-style-type: none"> - intra-abdominal free fluid - suspicion organ injury - suspicion retroperitoneal injury 3. Dislocated pelvic ring fracture and/or dislocated acetabulum fracture 4. Clinical suspicion of intraabdominal injury at physical examination 5. Subjective judgment of severity of injury by trauma leader <ul style="list-style-type: none"> - combined thoracic and pelvic injury - 'seatbelt sign' - chance fracture <p>X-thoracic and lumbar spine</p> <p>Not indicated when chest or abdominal CT is performed (reconstructions can be made)</p> <ol style="list-style-type: none"> 1. Complaints of the thoracic and lumbar spine 2. Tenderness of the thoracic and lumbar spine in the midline 3. Loss of consciousness 4. Deficits in peripheral neurologic examination 5. Painful distracting injuries <p>Pelvic CT (with iv contrast)</p> <ol style="list-style-type: none"> 1. All pelvic ring and acetabulum fractures unless conventional imaging is sufficient for adequate diagnosis and treatment 2. After reposition of hip luxation with suspicion of femoral head fractures and/or acetabulum fracture. <p>When CT-abdomen is performed, CT-pelvis is not necessary.</p> <p>Retrograde urethrogram</p> <ol style="list-style-type: none"> 1. Male patient with severe pelvic injury (type B and C) 2. Bleeding from the meatus, perineal injury or injury of the outer genital organs 3. Penetrating abdominal injury 4. In women only selectively after inspection <p>Imaging of the extremities</p> <p>When fractures/dislocations are suspected: conventional imaging and selective CT</p>
--	--

Appendix 3

Table 5 Follow-up for incidental findings

Characteristic	Total-body CT	Standard work-up	<i>p</i> *
Incidental finding of <i>major</i> relevance, <i>n</i>	24	9	0.129‡
Follow-up, <i>n</i> (%)	12 (50.0)	5 (55.6)	
No follow-up, <i>n</i> (%)	9 (37.5)	0 (0.0)	
Deceased in-hospital, <i>n</i> (%)*	3 (12.5)	4 (44.4)	
Incidental finding of <i>moderate</i> relevance, <i>n</i>	160	115	0.141†
Follow-up, <i>n</i> (%)	10 (6.3)	13 (11.3)	
No follow-up, <i>n</i> (%)	126 (78.8)	86 (74.8)	
Deceased in-hospital, <i>n</i> (%)*	24 (15.0)	16 (13.9)	
Incidental finding of <i>minor</i> relevance, <i>n</i>	257	166	0.735†
Follow-up, <i>n</i> (%)	1 (0.4)	1 (0.6)	
No follow-up, <i>n</i> (%)	227 (88.3)	141 (84.9)	
Deceased in-hospital, <i>n</i> (%)*	29 (11.3)	24 (14.5)	

* Incidental findings in patients who died in-hospital were excluded from this analysis

† Chi-square, ‡ Fisher's exact

Appendix 4

Table 6 Documentation of incidental findings

Characteristic	Total-body CT	Standard work-up	<i>p</i>
Incidental finding of <i>major</i> relevance, <i>n</i>	24	9	
Trauma letter, <i>n</i> (%)	10 (41.7)	5 (55.6)	0.697†
Discharge letter, <i>n</i> (%)	10 (41.7)	3 (33.3)	0.999†
Incidental finding of <i>moderate</i> relevance, <i>n</i>	160	115	
Trauma letter, <i>n</i> (%)	31 (19.4)	21 (18.3)	0.816*
Discharge letter, <i>n</i> (%)	20 (12.5)	18 (15.7)	0.455*
Incidental finding of <i>minor</i> relevance, <i>n</i>	257	166	
Trauma letter, <i>n</i> (%)	11 (4.3)	9 (5.4)	0.642†
Discharge letter, <i>n</i> (%)	13 (5.1)	6 (3.6)	0.632†

* Chi-square, † Fisher's exact

Appendix 5

Table 7 List of incidental findings in 1083 trauma patients, categorized by body region and clinical relevance

Location		Frequency	Percentage of total findings
Head	Major	mass, brain	4 0.5 %
	Moderate	aneurysm, brain, <5.5 cm	1 0.1 %
		cranial osteoma	5 0.7 %
		leukoaraiosis <50 years	2 0.3 %
		Minor	retention cyst
		leukoaraiosis >50 years	25 3.4 %
		brain calcification	16 2.2 %
		arachnoid cyst	10 1.4 %
		brain cyst	2 0.3 %
		cisterna magna, large	1 0.1 %
		colloid cyst	1 0.1 %
		parotid stone	1 0.1 %
	Neck	Major	-
Moderate		thyroid nodule	22 3.0 %
		cervical lymphadenopathy	3 0.4 %
		thyroid lesion	6 0.8 %
		goitre/struma	5 0.7 %
		mass, thyroid	1 0.1 %
		oesophageal hyperplasia	1 0.1 %
		thyroid cyst, complicated	1 0.1 %
Minor		cervical lymphadenopathy	5 0.7 %
		thyroid calcification	2 0.3 %
	thyroid cyst, simple	1 0.1 %	
Thorax	Major	pulmonary nodule, tumorous aspect	6 0.8 %
		mass, breast	1 0.1 %
		penetrating aortic ulcer	1 0.1 %
		pulmonary lesion, tumorous aspect	1 0.1 %
		thoracic lymphadenopathy	1 0.1 %
	Moderate	cardiomegaly	10 1.4 %
		pulmonary nodule, relevant	6 0.8 %
		COPD	3 0.4 %
		aneurysm, thoracic, <5,5 cm	1 0.1 %
		atelectasis	2 0.3 %
		gynecomastia	1 0.1 %

Table 7 (continued)

Location	Frequency	Percentage of total findings
cutaneous nodule, relevant	1	0.1 %
diaphragmatic hernia	3	0.4 %
hepatic cyst, complicated	4	0.5 %
hepatic lesion, relevant	6	0.8 %
hernia, inguinal	1	0.1 %
hernia, umbilical	1	0.1 %
horseshoe kidney	3	0.4 %
hydrocele testis	1	0.1 %
hydronephrosis	4	0.5 %
mono-kidney	1	0.1 %
pancreatic atrophy	7	1.0 %
pancreatic calcification	3	0.4 %
pancreatic cyst	3	0.4 %
pancreatic steatosis	2	0.3 %
pneumaturia	1	0.1 %
porcelain gallbladder	1	0.1 %
prostatic hypertrophy	8	1.1 %
renal atrophy	2	0.3 %
renal lesion, complicated	1	0.1 %
Riedel's hepatic lobe	1	0.1 %
splenic lesion, relevant	2	0.3 %
suspected fibromuscular dysplasia	1	0.1 %
vesical calculus	1	0.1 %
Minor		
renal cyst, simple	97	13.3 %
hepatic cyst, simple	50	6.8 %
diverticulosis	35	4.8 %
spleen, accessory	20	2.7 %
hepatic lesion, simple	11	1.5 %
abdominal lymphadenopathy	6	0.8 %
adrenal cyst, simple	2	0.3 %
adrenal lesion, simple	4	0.5 %
adrenal nodule, simple	3	0.4 %
bladder diverticulum	3	0.4 %
corpus luteum cyst	1	0.1 %
duplex collecting system	1	0.1 %
fluid in rectouterine pouch	3	0.4 %
intestinal malrotation	1	0.1 %
ovarian cyst, <5 cm	3	0.4 %
prostatic calcification	10	1.4 %
renal calcification	1	0.1 %
renal cortex, thinning	2	0.3 %
renal ectopia	1	0.1 %
renal lesion, simple	3	0.4 %
splenic cyst, simple	4	0.5 %
urachal cyst	1	0.1 %
uterine calcification	4	0.5 %

Table 7 (continued)

Location		Frequency	Percentage of total findings		
Extremities	Major	uterine fibroid	1	0.1 %	
		-			
	Moderate	bone lytic lesion	2	0.3 %	
		Minor	bone cyst, simple	1	0.1 %
			bone lesion, simple	1	0.1 %

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Asha S, Curtis KA, Grant N et al (2012) Comparison of radiation exposure of trauma patients from diagnostic radiology procedures before and after the introduction of a panscan protocol. *Emerg Med Australas* 24:43–51
- Sierink JC, Saltzherr TP, Wirtz MR, Streekstra GJ, Beenen LF, Goslings JC (2013) Radiation exposure before and after the introduction of a dedicated total-body CT protocol in multitrauma patients. *Emerg Radiol* 20:507–512
- Sierink JC, Treskes K, Edwards MJ et al (2016) Immediate total-body CT scanning versus conventional imaging and selective CT scanning in patients with severe trauma (REACT-2): a randomised controlled trial. *Lancet*. doi:10.1016/s0140-6736(16)30932-1
- Paluska TR, Sise MJ, Sack DI, Sise CB, Egan MC, Biondi M (2007) Incidental CT findings in trauma patients: incidence and implications for care of the injured. *J Trauma* 62:157–161
- Munk MD, Peitzman AB, Hostler DP, Wolfson AB (2010) Frequency and follow-up of incidental findings on trauma computed tomography scans: experience at a level one trauma center. *J Emerg Med* 38:346–350
- Barrett TW, Schierling M, Zhou C et al (2009) Prevalence of incidental findings in trauma patients detected by computed tomography imaging. *Am J Emerg Med* 27:428–435
- Hoffstetter P, Herold T, Daneschnejad M et al (2008) Non-trauma-associated additional findings in whole-body CT examinations in patients with multiple trauma. *Rofo* 180:120–126
- Sierink JC, Saltzherr TP, Russchen MJ et al (2014) Incidental findings on total-body CT scans in trauma patients. *Injury* 45:840–844
- Sierink JC, Saltzherr TP, Beenen LF et al (2012) A multicenter, randomized controlled trial of immediate total-body CT scanning in trauma patients (REACT-2). *BMC Emerg Med* 12:4
- Beenen LF, Sierink JC, Kolkman S et al (2015) Split bolus technique in polytrauma: a prospective study on scan protocols for trauma analysis. *Acta Radiol* 56:873–880
- Cheng T, Dumire R, Golden S, Gregory J (2013) Impact on patient care of discordance in radiology readings between external overnight radiology services and staff radiology readings at a level 1 trauma center. *Am J Surg* 205:280–282, discussion 282–283
- Ekeh AP, Walusimbi M, Brigham E, Woods RJ, McCarthy MC (2010) The prevalence of incidental findings on abdominal computed tomography scans of trauma patients. *J Emerg Med* 38:484–489
- Lumbreras B, Donat L, Hernandez-Aguado I (2010) Incidental findings in imaging diagnostic tests: a systematic review. *Br J Radiol* 83:276–289
- van Vugt R, Dekker HM, Deunk J et al (2011) Incidental Findings on Routine Thoracoabdominal Computed Tomography in Blunt Trauma Patients. *J Trauma*. doi:10.1097/TA.0b013e3182166b4b
- Bosniak MA (1986) The current radiological approach to renal cysts. *Radiology* 158:1–10
- MacMahon H, Austin JH, Gamsu G et al (2005) Guidelines for management of small pulmonary nodules detected on CT scans: a statement from the Fleischner Society. *Radiology* 237:395–400
- Devine AS, Jackson CS, Lyons L, Mason JD (2010) Frequency of incidental findings on computed tomography of trauma patients. *West J Emerg Med* 11:24–27
- Messersmith WA, Brown DF, Barry MJ (2001) The prevalence and implications of incidental findings on ED abdominal CT scans. *Am J Emerg Med* 19:479–481
- Thompson RJ, Wojcik SM, Grant WD, Ko PY (2011) Incidental Findings on CT Scans in the Emergency Department. *Emerg Med Int* 2011:624847
- Mets OM, de Jong PA, Chung K, Lammers JJ, van Ginneken B, Schaefer-Prokop CM (2016) Fleischner recommendations for the management of subsolid pulmonary nodules: high awareness but limited conformance - a survey study. *Eur Radiol*. doi:10.1007/s00330-016-4249-y